

(1)  $\begin{pmatrix} 54 & 110 \\ -3 & -11 \end{pmatrix}$  (2)  $\frac{1}{66} \begin{pmatrix} -54 & -110 \\ 3 & 11 \end{pmatrix}$

(3)  $\frac{1}{66} \begin{pmatrix} 54 & 110 \\ 3 & 11 \end{pmatrix}$  (4) none of these

5. The value of the determinant  $\begin{vmatrix} \cos \alpha & \sin \alpha & 0 \\ -\sin \alpha & \cos \alpha & 0 \\ 0 & 0 & 1 \end{vmatrix}$  is :

(1) -1 (2) 0 (3) 1 (4)  $\cos 2\alpha$

6. If  $\omega$  is the cube root of unity then the value of  $\begin{vmatrix} 1 & \omega & \omega^2 \\ \omega & \omega^2 & 1 \\ \omega^2 & 1 & \omega \end{vmatrix}$  will be :

(1)  $\omega$  (2)  $\omega^2 + 1$  (3) 0 (4) 1

7. If  $(1+x)^n = C_0 + C_1x + C_2x^2 + \dots + C_nx^n$ , then  $C_0 + \frac{C_1}{2} + \frac{C_2}{3} + \dots + \frac{C_n}{n+1}$  is equal to :

(1)  $\frac{2^n - 1}{n+1}$  (2)  $\frac{2^n - 1}{n-1}$  (3)  $\frac{2^{n-1} - 1}{n+1}$  (4)  $\frac{2^{n+1} - 1}{n+1}$

8. If  $(1+x)^n = C_0 + C_1x + C_2x^2 + \dots + C_nx^n$ , then  $C_0C_2 + C_1C_3 + C_2C_4 + \dots = C_{n-2} C_n$  is equal to :

(1)  $\frac{2n}{n-2}$  (2)  $\frac{2n}{n}$  (3)  $\frac{2n}{n-2}$  (4) none of these

9. If the ratio of the second and third term in the expansion of  $(a+b)^2$  is equal to the ratio of third and fourth term in the expansion of  $(a+b)^{n+3}$  then the value of n is equal to :

(1) 6 (2) 4 (3) 5 (4) 3

10. The number of different words that can be formed by using the letters of the word 'MISSISSIPPI' is :

(1) 5067 (2) 6705 (3) 1520 (4) 2520

11. If  $a^2, b^2, c^2$  are in A.P. then  $\frac{1}{b+c^2}, \frac{1}{c+a}, \frac{1}{a+b}$  will be :

(1) in H.P. (2) in G.P. (3) in A.P. (4) in arithmetic progression

12. A bag contains 5 white and 3 black balls. Two balls are drawn at random then the probability that out of the two one ball is red and other is black will be :

- (1)  $\frac{15}{56}$       (2)  $\frac{11}{28}$       (3)  $\frac{15}{28}$       (4)  $\frac{2}{7}$

13. Two dice are thrown together then the probability that the sum of numbers appearing on the dice is 7 :

- (1)  $\frac{1}{6}$       (2)  $\frac{1}{12}$       (3)  $\frac{5}{36}$       (4) none of these

14. The position vectors of the two points A and B are a and b respectively then the position vector of the point C which divides AB in the ration 2 : 1 will be :

- (1)  $\frac{1+b}{3}$       (2)  $\frac{2a+b}{3}$       (3)  $\frac{a+2b}{3}$       (4) none of these

15. If  $a + b = a - b$ , then the angle between a and b will be :

- (1)  $180^0$       (2)  $90^0$       (3)  $60^0$       (4)  $0^0$

16. The area of the region bounded by the curve  $y = \sin^2 x$ , x-axis and the lines  $x = 0$ ,  $x = \pi/2$  is

- (1)  $\pi$       (2)  $\pi/8$       (3)  $\frac{\pi}{4}$       (4)  $\frac{\pi}{2}$

17.  $\int_0^{\pi/2} \frac{\sin x}{\sqrt{1 + \sin 2x}} dx$  is equal to :

- (1)  $\pi$       (2)  $2\pi$       (3)  $\frac{\pi}{4}$       (4)  $\frac{\pi}{2}$

18.  $\int xe^x dx$  is equal to :

- (1)  $(x - 1)e^x + C$       (2)  $(1 - x)e^x + C$   
 (3)  $(1 - x)e^x + C$       (4) none of these

19.  $\int x^2 \sin x^3 dx$  is equal to :

- (1)  $\frac{1}{3} \sin x^3 + C$       (2)  $-\frac{1}{3} \sin x^3 + C$   
 (3)  $\frac{1}{3} \cos x^3 + C$       (4)  $-\frac{1}{3} \cos x^3 + C$

20. The max. value of  $\sin x + \cos x$  :

- (1) 1      (2)  $\frac{1}{\sqrt{2}}$       (3)  $\sqrt{2}$       (4) none of these

21. The angle between the curves  $y = x$  and  $y^2 = 4x$  at origin will be :

- (1) 0                      (2)  $\frac{\pi}{3}$                       (3)  $\frac{\pi}{4}$                       (4) none of these

**22. If the volume of a balloon is increasing at the rate of 25 cm<sup>3</sup>/sec., then if the radius of the balloon is 5 cm. then the rate of change of the surface area is :**

- (1) 20 cm<sup>2</sup>/sec.                      (2) 10 cm<sup>2</sup>/sec.                      (3) 5 cm<sup>2</sup>/sec.                      (4) 10 cm./sec.

**23. The differential coefficient of x<sup>x</sup> is :**

- (1) x<sup>x</sup> log<sub>e</sub> x                      (2) x<sup>x</sup> (1 + log<sub>e</sub> x)                      (3) x<sup>x</sup> (1 - log<sub>e</sub> x)                      (4) none of these

**24.  $\frac{d}{dx} (\sin x)^{\tan x}$  is equal to :**

- (1)  $(\sin x)^{\tan x} [1 - \sec^2 x \log \sin x]$   
 (2)  $(\tan x)^{\sin x} \log \sec^2 x \log \sin x]$   
 (3)  $(\sin x)^{\tan x} [1 + \sec^2 x \log \sin x]$   
 (4) none of these

**25. If  $y = \sec^{-1} \left( \frac{x+1}{x-1} \right) + \sin^{-1} \left( \frac{x-1}{x+1} \right)$**

**then  $\frac{dy}{dx}$  is equal to :**

- (1)  $\infty$                       (2) 0                      (3) 1                      (4) -1

**26. If  $y = \sin^{-1} \left( \frac{2x}{1+x^2} \right)$ , then  $\frac{dy}{dx}$  is equal to :**

- (1)  $-\frac{1}{1+x^2}$                       (2)  $-\frac{2}{1+x^2}$   
 (3)  $\frac{2}{1+x^2}$                       (4)  $\frac{2x}{1+x^2}$

**27. If  $x\sqrt{1+y} + y\sqrt{1+x} = 0$  then  $\frac{dy}{dx}$  is equal to :**

- (1)  $\frac{1}{1+x^2}$                       (2)  $-\frac{1}{1+x^2}$   
 (3)  $\frac{2}{(1+x)^2}$                       (4) none of these

**28. The continuous product of the roots of (-1)<sup>2/3</sup> is :**

- (1)  $\omega^2$                       (2)  $\omega$                       (3) 0                      (4) 1

**29. The value of  $\sin^{-1} x$  :**

- (1)  $\log(x - \sqrt{x^2 - 1})$                       (2)  $\log(x + \sqrt{x^2 - 1})$

(3)  $\log(x + \sqrt{x^2 + 1})$

(4)  $\frac{1}{2} \log \frac{1+x}{1-x}$

**30. The equation of a st-line passing through the point (1,2) and making equal angles to with axes, will be :**

(1)  $x - y - 2 = 0$

(2)  $x + y + 1 = 0$

(3)  $x - y = 1$

(4)  $x + y = 1$

**31. If the vertices of a parallelogram are (0,0), (2,1), (1,3) and (1,2) then the angle between their diagonals will be:**

(1)  $\frac{\pi}{4}$

(2)  $\frac{3\pi}{2}$

(3)  $\frac{\pi}{2}$

(4)  $\frac{\pi}{3}$

**32. The equation of line which is parallel to the straight line  $3x + 4y - 7 = 0$  and passing through (1,2) is :**

(1)  $3x + 4y = 11$

(2)  $3x + 4y + 11 = 0$

(3)  $4x - 3y + 2 = 0$

(4)  $3x + 4y + 7 = 0$

**33. The pole of the straight line  $9x + y - 28 = 0$  w.r.t. the circle  $x^2 + y^2 = 16$  will be:**

(1)  $\left(\frac{33}{7}, \frac{3}{7}\right)$

(2)  $\left(\frac{33}{7}, \frac{4}{7}\right)$

(3)  $\left(\frac{4}{7}, \frac{36}{7}\right)$

(4)  $\left(\frac{36}{7}, \frac{4}{7}\right)$

**34. The equation of the tangent from origin to the circle  $x^2 + y^2 - 2rx - 2hy + h^2 = 0$  is:**

(1)  $(h^2 - r^2)x + 2rhy = 0$

(2)  $y = 0$

(3)  $x - y = 0$

(4)  $(h^2 - r^2)x - 2rhy = 0$

**35. If a tangent at a point p to the parabola meets to the directrix at Q. If S is the focus of the parabola then  $\angle PSQ$  is equal to :**

(1)  $\pi$

(2)  $\frac{\pi}{2}$

(3)  $\frac{\pi}{3}$

(4)  $\frac{\pi}{4}$

**36. If  $f(y) = \log y$ , then  $f(y) + f(1/y)$  is equal to :**

(1) 2

(2) 0

(3) -1

(4) 1

**37.  $\lim_{x \rightarrow 0} \frac{e^x - \log(1+x)}{x^2}$  is equal to :**

(1)  $\frac{1}{2}$

(2)  $\frac{1}{3}$

(3)  $\frac{3}{2}$

(4)  $\frac{2}{3}$

38. If  $\alpha$  and  $\beta$  are the roots of the equation  $1 - (1 + n^2 + n^4) = 0$  then  $\alpha^2 + \beta^2$  is equal to :

- (1)  $2n^2$       (2)  $n^2$       (3)  $-n^2$       (4)  $n^2 + 2$

39. The H.M. between 1 and  $\frac{1}{16}$  will be :

- (1)  $\frac{17}{2}$       (2)  $\frac{2}{17}$       (3)  $\frac{17}{32}$       (4)  $\frac{32}{17}$

40. If for two numbers G.M. is 4 and A.M. is 5, then H.M. will be :

- (1)  $\frac{25}{15}$       (2)  $\frac{17}{8}$       (3)  $\frac{16}{5}$       (4)  $\frac{5}{16}$

41. If  ${}^{10}C_r = {}^{10}C_{r+2}$  then  ${}^5C_r$  is equal to :

- (1) 360      (2) 120      (3) 10      (4) 5

42. The value of  $1 + \frac{1}{4} + \frac{1.3}{4.8} + \frac{1.3.5}{4.8.12} + \dots$  is :

- (1)  $\sqrt{3/2}$       (2)  $\sqrt{2}$       (3) 2      (4)  $3/2$

43. If  $(1+x)^n = C_0 + C_1x + C_2x^2 + \dots + C_nx^n$  then  $\frac{C_1}{C_0} + \frac{2C_2}{C_1} + \frac{3C_3}{C_2} + \dots + \frac{nC_n}{C_{n-1}}$  is equal to :

- (1)  $\frac{n(n+1)}{2}$       (2)  $\frac{n(n^2+1)}{2}$       (3)  $\frac{n(n+1)}{n!}$       (4)  $\frac{n(n-1)}{2}$

44. In the expansion of  $\left(2^4 - \frac{1}{x^7}\right)$  the term independent of x is :

- (1) -32190      (2) 114050      (3) 42240      (4) 330

45. The value of the determinant  $\begin{vmatrix} 4 & -6 & 1 \\ -1 & -1 & 1 \\ -4 & 11 & -1 \end{vmatrix}$  is :

- (1) 0      (2) -25      (3) 25      (4) none of these

46. If  $\begin{vmatrix} 1 & 2 & 4 \\ 3 & 6+x & 7 \end{vmatrix} = 0$ , then the value of x will be :

5 10 4+x

- (1) 3                      (2) 0                      (3) 1                      (4) none of these

47. If  $A = \begin{pmatrix} d_1 & 0 & 0 \\ 0 & d_2 & 0 \\ 0 & 0 & d_3 \end{pmatrix}$  then  $\text{adj } A =$

(1)  $\begin{pmatrix} d_1^{-1} & 0 & 0 \\ 0 & d_2^{-1} & 0 \\ 0 & 0 & d_3^{-1} \end{pmatrix}$

(2)  $\begin{pmatrix} d_2 d_3 & 0 & 0 \\ 0 & d_1 d_3 & 0 \\ 0 & 0 & d_1 d_2 \end{pmatrix}$

(3)  $\begin{pmatrix} d_2 d_2 & 0 & 0 \\ 0 & d_1 d_3 & 0 \\ 0 & 0 & d_1 d_3 \end{pmatrix}$

(4)  $\begin{pmatrix} d_1 d_3 & 0 & 0 \\ 0 & d_2 d_3 & 0 \\ 0 & 0 & d_1 d_2 \end{pmatrix}$

48. If  $A = \begin{pmatrix} 2 & 4 \\ 0 & 3 \end{pmatrix}$  and  $B = \begin{pmatrix} 1 & 2 \\ 0 & 5 \end{pmatrix}$ , then  $4A - 3B$  is equal to :

(1)  $\begin{pmatrix} 1 & 2 \\ 0 & 2 \end{pmatrix}$                       (2)  $\begin{pmatrix} -5 & -10 \\ 0 & 3 \end{pmatrix}$                       (3)  $B = \begin{pmatrix} 5 & 10 \\ 0 & -3 \end{pmatrix}$                       (4)  $A = \begin{pmatrix} 7 & 14 \\ 0 & 7 \end{pmatrix}$

49. If  $A = \begin{pmatrix} \cos x & \sin x \\ -\sin x & \cos x \end{pmatrix}$ , then  $A^{-1}$  is equal to :

- (1)  $\begin{pmatrix} \cos x & \sin x \\ \sin x & \cos x \end{pmatrix}$  (2)  $\begin{pmatrix} \cos x & -\sin x \\ \sin x & \cos x \end{pmatrix}$  (3)  $\begin{pmatrix} \cos x & \sin x \\ -\sin x & \cos x \end{pmatrix}$  (4) none of these

50. A card is drawn at random from a pack of playing cards. The probability that it is red or an ace, is :

- (1)  $\frac{1}{13}$  (2)  $\frac{1}{52}$  (3)  $\frac{17}{52}$  (4)  $\frac{4}{13}$

51. If the sum of two unit vector is also a unit vector then the magnitude of their difference will be :

- (1) 1 (2)  $\sqrt{3}$  (3)  $\frac{1}{\sqrt{3}}$  (4)  $\sqrt{2}$

52. The unit vector perpendicular to the vectors  $6\mathbf{i} + 2\mathbf{j} + 3\mathbf{k}$  and  $3\mathbf{i} - 6\mathbf{j} - 2\mathbf{k}$  will be:

- (1)  $\frac{2\mathbf{i} - 3\mathbf{j} - 6\mathbf{k}}{7}$  (2)  $\frac{2\mathbf{i} - 3\mathbf{j} + 6\mathbf{k}}{7}$  (3)  $\frac{2\mathbf{i} + 3\mathbf{j} - 6\mathbf{k}}{7}$  (4)  $\frac{2\mathbf{i} + 3\mathbf{j} + 6\mathbf{k}}{7}$

53. The area of the region bounded by the curves  $y^2 = 4ax$ ,  $x = 0$  and  $x = a$  is

- (1)  $4\pi a^2$  (2)  $3\pi a^2$  (3)  $2\pi a^2$  (4)  $\pi a^2$

54. The area of the region bounded by the curves  $y^2 = 4ax$ ,  $x = 0$  and  $x = a$  is :

- (1)  $\frac{5}{3} a^2$  (2)  $\frac{2}{3} a^2$  (3)  $\frac{8}{3} a^2$  (4)  $\frac{4}{3} a^2$

55.  $\int \cos^3 x \, dx$  is equal to

- (1)  $\frac{\sin 3x}{4} + 3 \sin x + C$   
 (2)  $\frac{\sin 3x}{3} + \frac{\sin x}{2} + C$   
 (3)  $\sin 3x + C$   
 (4)  $\frac{\sin 3x}{12} + \frac{3}{4} \sin x + C$

56. If  $x = a(t + \sin t)$  and  $y = a(1 - \cos t)$  then  $\frac{dy}{dx}$  is equal to :

- (1)  $\tan t$  (2)  $\tan 2t$  (3)  $\cot(t/2)$  (4)  $\tan(t/2)$

57. If  $x = t^2$  and  $y = 2t$ , then the normal at  $t = 1$  is :

- (1)  $x + y - 3 = 0$       (2)  $x + y - 1 = 0$   
 (3)  $x + y + 1 = 0$       (4)  $x + y + 3 = 0$

58.  $f(x) = 2x^3 - 9x^2 + 12x + 29$  is a monotonic decreasing function when :

- (1)  $1 < x < 2$     (2)  $x > 1$       (3)  $x > 2$       (4)  $x < 2$

59. The height of the cylinder of maximum volume that can be inscribed in a sphere of radius  $r$  is :

- (1)  $2\sqrt{3}r$       (2)  $\frac{2r}{\sqrt{3}}$       (3)  $r\sqrt{3}$       (4)  $\frac{r}{\sqrt{3}}$

60.  $\int \sec x \, dx$  is equal to :

- (1)  $\log \sin x + C$   
 (2)  $\log \tan (x/2) + C$   
 (3)  $-\log (\sec x - \tan x) + C$   
 (4)  $\log \tan \left[ \frac{\pi}{2} + \frac{\pi}{4} \right] + C$

61. The differential coefficient of  $\sin^{-1} \left( \frac{1-x^2}{1+x^2} \right)$  w.r.t.  $x$  is :

- (1)  $-\frac{2}{1+x^2}$     (2)  $\frac{2}{1+x^2}$       (3)  $\frac{1}{1+x^2}$     (4) none of these

62.  $d(\sec^{-1} x)$  is equal to :

- (1)  $\frac{1}{x\sqrt{x-1}}$       (2)  $\frac{1}{x\sqrt{x^2-1}}$       (3)  $\frac{1}{x\sqrt{1+x^2}}$       (4)  $\frac{1}{x\sqrt{1-x^2}}$

63. The differential coefficient of  $\tan^{-1} \sqrt{\frac{1-x^2}{1+x^2}}$  w.r.t.  $x$  is :

- (1)  $\frac{1}{2}$       (2)  $1$       (3)  $-\frac{1}{2}$       (4) none of these

64.  $\lim_{x \rightarrow \frac{\pi}{2}} \frac{\tan 2x - x}{3x - \sin x}$  is equal to :

- (1)  $0$       (2)  $1$       (3)  $\frac{1}{3}$       (4)  $\frac{1}{4}$

65. The differential coefficient of  $\sin^{-1} x$  w.r.t.  $\cos^{-1} \sqrt{1-x^2}$  is :

- (1)  $-\frac{1}{\sqrt{1+x^2}}$       (2)  $\frac{1}{\sqrt{1-x^2}}$       (3)  $\frac{2}{\sqrt{1-x^3}}$       (4) none of these

66. The sum of 20 terms of the series  $1 + 4 + 7 + 10 + \dots$  is :



- (1) 290      (2) 490      (3) 590      (4) none of these

67. Which terms of the series  $\frac{1}{2}, \frac{-1}{2}, 1-2, \dots$  is  $-128$  :

- (1)  $10^{\text{th}}$       (2)  $8^{\text{th}}$       (3)  $9^{\text{th}}$       (4)  $12^{\text{th}}$

68. If  ${}^n P_4 : {}^n P_5 = 1 : 2$ , then  $n$  is equal to :

- (1) 2      (2) 4      (3) 5      (4) 6

69.  $\frac{(\cos 3\theta + i \sin 3\theta)^5 (\cos 2\theta + i \sin 2\theta)^4}{(\cos \theta + i \sin \theta)^4 (\cos \theta + i \sin \theta)^6}$  is equal to :

- (1)  $\cos 27\theta - i \sin 27\theta$   
 (2)  $\cos 33\theta - i \sin 33\theta$   
 (3)  $\cos 33\theta + i \sin 33\theta$   
 (4)  $\cos 27\theta + i \sin 27\theta$

70. The value of  $\cos^{-1} x$  is :

- (1)  $\log(x - \sqrt{x^2 - 1})$       (2)  $\log(x + \sqrt{x^2 - 1})$   
 (3)  $\log(x + \sqrt{x^2 + 1})$       (4)  $\log(x - \sqrt{x^2 + 1})$

71. Find the equation of the straight line which is perpendicular to the line  $\frac{x}{a} - \frac{y}{b} = 1$  and passes through the point where the given st-line cuts the x-axis :

- (1)  $ax + by = a^2$       (2)  $ax - by = a^2$       (3)  $ax + by = b^2$       (4)  $bx - ay = ab$

72. If the lines  $x + y = 1$ ,  $2x - y = 0$  and  $x + 2y + \lambda = 0$  are concurrent then  $\lambda$  is equal to :

- (1)  $-\frac{2}{3}$       (2)  $\frac{2}{3}$       (3)  $-\frac{5}{3}$       (4)  $\frac{5}{3}$

73. If two vertices of a triangle are (6,4), (2,6) and its centroid is (4, 6) then its third vertex will be :

- (1) (6,4)      (2) (8,4)      (3) (4,8)      (4) none of these

74. The radical axis of the circles  $2x^2 + 2y^2 - 7x = 0$  and  $x^2 + y^2 - 4y - 7 = 0$  is :

- (1)  $8x - 7y + 14 = 0$       (2)  $7x - 8y + 14 = 0$   
 (3)  $7x - 8y - 14 = 0$       (4)  $7x + 8y + 14 = 0$

75. The equation of the polar line w.r.t. the pole (1, -2) to the circle  $x^2 + y^2 - 2x - 6y + 5 = 0$  is :

- (1)  $x + y - 1 = 0$       (2)  $x + y + 1 = 0$       (3)  $y = 2$       (4)  $x = 2$

76. The vertex of the parabola  $x^2 - y + 6x + 10 = 0$  is :

- (1) (3,1)                      (2) (3, -1)                      (3) (3,-2)                      (4) (-3, 1)

77. If  $f(\theta) = \tan \theta$  then the value of  $\frac{f(\theta) - f(\phi)}{1 + f(\theta)f(\phi)}$  is :

- (1)  $\theta - \phi$                       (2)  $f(\theta/\phi)$                       (3)  $f(\theta - \phi)$                       (4)  $f(\theta + \phi)$

78.  $\lim_{x \rightarrow 0} \frac{x^2 - 3x + 2}{2x^2 + x - 3}$  is equal to :

- (1) 0                      (2) 2                      (3)  $\frac{1}{2}$                       (4)  $\infty$

79.  $\lim_{x \rightarrow 0} \left[ \frac{\sqrt{1+x} - \sqrt{1-x}}{x} \right]$  is equal to :

- (1) -1                      (2) 1                      (3) 2                      (4)  $\frac{1}{2}$

80. The equation of the normal at a point of intersection of line  $2x + y = 3$  and curve  $yx^2 + y^2 = 5$  is :

- (1)  $2x + 2y + 3 = 0$                       (2)  $x - y + 4 = 0$                       (3)  $x - 4y + 3 = 0$                       (4)  $x + y + 2 = 0$

81. If  $f(x) = \frac{x - 3}{x + 1}$ , then  $f[f(f(x))]$  is equal to :

- (1)  $\frac{-1}{x}$                       (2)  $-x$                       (3)  $\frac{1}{x}$                       (4)  $x$

82. The modulus of  $\frac{1+i}{1-i}$  is :

- (1)  $\sqrt{2}$                       (2) 2                      (3)  $\frac{1}{2}$                       (4) 1

83. The value of  $\frac{4\sqrt{\sqrt{3}}}{7} - \frac{\sqrt{\sqrt{3}}}{7}$  is

- (1)  $\frac{3\sqrt{3}}{7}$                       (2)  $-\frac{3\sqrt{3}}{7}$                       (3)  $\frac{3\sqrt{3}}{7} i$                       (4) none of these

84.  $\frac{1-2i}{2+i} + \frac{4-i}{3+2i}$  is equal to :

- (1)  $\frac{10}{13} + \frac{24}{13} i$                       (2)  $\frac{10}{13} - \frac{24}{13} i$                       (3)  $\frac{24}{13} - \frac{10}{13} i$                       (4)  $\frac{24}{13} + \frac{10}{13} i$

85. If  $z = 5 + 3i$  then the value of  $|z - 2|$  will be :

- (1)  $\sqrt{13}$                       (2)  $2\sqrt{3}$                       (3)  $3\sqrt{2}$                       (4) 13

86. The imaginary part of  $\frac{1-i}{1+i}$  is :

- (1)  $-i$             (2)  $-1$             (3)  $1$             (4)  $i$

87. If  $z_1 = 1 + 2i$  and  $z_2 = i - 1$ , then  $z_1/z_2$  is equal to :

- (1)  $\frac{1}{2} - \frac{3}{2}i$             (2)  $-\frac{1}{2} + \frac{3}{2}$             (3)  $\frac{1}{2} - \frac{3}{2}i$             (4) none of these

88. The amplitude of  $1 - \sqrt{3}i$  is :

- (1)  $\frac{-2\pi}{3}$             (2)  $\frac{-\pi}{3}$             (3)  $\frac{2\pi}{3}$             (4)  $\frac{\pi}{3}$

89. If  $\alpha$  and  $\beta$  are the roots of the equation  $x^2 + px + q = 0$  then the value of  $\alpha^3 + \beta^3 = 0$  will be :

- (1)  $p^3 - 3pq$             (2)  $-(p^3 + 3pq)$             (3)  $p^3 + 3pq$             (4)  $-p^3 + 3pq$

90. If  $\alpha$  and  $\beta$  are the roots of the equation whose roots are  $\frac{1}{\alpha}, \frac{1}{\beta}$  is :

- (1)  $x^2 + x + 1 = 0$             (2)  $x^2 - x + 1 = 0$             (3)  $x^2 - x = 1$             (4)  $x^2 - x = 1$

91. If  $z = \frac{(1+i)(2+i)}{3+i}$  then  $|z|$  is equal to :

- (1)  $-\frac{1}{2}$             (2)  $\frac{1}{2}$             (3)  $1$             (4)  $-1$

92. The slope of the tangent to the parabola  $y^2 = 4ax$  point  $(at^2, 2at)$  will be :

- (1)  $-t$             (2)  $-1/t$             (3)  $1/t$             (4)  $t$

93. If  $a = i - 2j$  and  $b = 2i + \lambda k$  are the parallel vectors then  $\lambda$  is equal to :

- (1)  $-2$             (2)  $2$             (3)  $-4$             (4)  $4$

94. A stone is thrown in silent water, the ripples are moving at the rate of 6 cm/sec. then the rate of change of the are when the radius of the unite is 10 cm. at the time when radius of the circle is 10 cm, then the rate at which its area increases is :

- (1)  $120 \text{ m}^2/\text{sec.}$             (2)  $\pi \text{ cm}^2/\text{sec.}$             (3)  $120 \text{ cm}^2/\text{sec.}$             (4)  $120 \pi \text{ cm}^2/\text{sec.}$

95. A dice is thrown then the probability that the sum of the number is 1 or 6 is :

- (1)  $1/6$             (2)  $1/3$             (3)  $2/3$             (4)  $3/4$

96. The value of  $\cos h(\pi i)$  is :

- (1)  $0$             (2)  $1$             (3)  $-1$             (4) none of these

97. For  $Z_1, Z_2 \in \mathbb{C}$  the value of  $|Z_1 + Z_2|^2 + |Z_1 - Z_2|^2$  will be :

- (1)  $2(|Z_1|^2 + |Z_2|^2)$             (2)  $|Z_1|^2 + |Z_2|^2$             (3)  $|Z_1|^2 + |Z_2|^2 - |Z_1| - |Z_2|$             (4)  $2(|Z_1|^2 - |Z_2|^2)$

