<u>-</u>	trains are approaching e $y_2 = a \sin 208 \pi t$	ach other. http://www.u	upadmission.com/	
A. 8	B. 4	C. 1	D. 0	
2. One of the geo-static A. New Delhi	onary satellites of India is B. Mumbai	s vertically above C. Allahabad	D. None of these	
3. Light of wavelength equal to	2400 x 10 ⁻¹⁰ m in air will	become light of waveler	ngth in glass ($\mu = 1.5$)	
A. 1600 x 10 ⁻¹⁰ m	B. 7200 x 10 ⁻¹⁰ m	C. 1080 x 10 ⁻¹⁰ m	D. none of these	
	ry to primary turns is 4:5 all losses) to power input	<u> </u>	at will be the ratio of	
A. 4:9	B. 9:4	C. 5:4	D. 1:1	
5. Lenz's law applies toA. electrostaticsC. electro-magnetic ind		B. lenses D. cinema slides		
	proton come close to each	n other and annihilate, ho	ow much energy will be	
A. 1.5 x 10 ⁻¹⁰ J	B. 3 x 10 ⁻¹⁰ J	C. 4.5 x 10 ⁻¹⁰ J	D. none of these	
7. If Sn is doped with A ?	s, what will be the result			
A. <i>n</i> -type B. <i>p</i> -type semi-conductor conductor	C. intrinsic semi-conductor D. none of these			
8. A charge is placed at faces?	the centre of a cube, wh	at is the electric flux pas	ssing through one of its	
A. $(1/6) \times (q/\epsilon_0)$	B. q/ϵ_0	C. $6q/\epsilon_0$	D. None of these	
9. What is the degree of A. 1	f freedom in case of a mo B. 3	ono atomic gas ? C. 5	D. None of these	
10. The ratio of secondary to primary turns is 4:5. If power input is P , what will be the ratio of power output (neglect all losses) to power input?				
Α.	В.	C.	D.	

11. Speed of recession of galaxy is proportional to its distance

		http://www.	upadmission.com/
A. directly	B. inversely	C. exponentially	D. none of these
12. If a substance goes	in a magnetic field and i	s pushed out of it, what i	is it ?
A. Paramagnetic	B. Ferromagnetic	C. Diamagnetic	D. Antiferromagnetic
13. Which is not a scale	ar quantity?		
A. Work	B. Power	C. Torque	D. Gravitational Constant
14. Minimum energy re	equired to excite an elect	ron in a Hydrogen atom	in ground state is:
A13.6 eV	B. 13.6 eV	C. 10.2 eV	D. 3.4 eV
15. If Gravitational Consatellite orbiting around	_	me, what will remain und	changed in case of a
A. Time period	B. Orbiting radius	C. Tangential velocity	D. Angular velocity
in front of one of the sl	its of Young's Double Sl	$\mu = 1.5$ and thickness $t =$ it experiment, how much slits is 5.0×10^{-3} cm and	h will be the shift in the
A. 5 cm	B. 2.5 cm	C. 0.25 cm	D. 0.1 cm
-	pagate in optical fibres?		
A. Total internal reflection	B. Refraction	C. Reflection	D. None of these
18. Dispersion of light	is due to		
A. wavelength 19. Which of the follow	B. intensity of light ving conclusions is corre	C. density of medium ct regarding	D. none of these
a stationary body? A. No force is acting or	n the hody		
•	s acing on the body is ze	ro	
C. The body is in vacuu	•		
D. The forces acting on	the body do not constitu	ıte a couple	
20. Energy released in	stars is due to		
A. Fission	B. Fusion	C. Combustion	D. Chemical reaction
21. 13 days is the half-1/16th of the original si	•	After how many days, the	e sample will become
A. 52	B. 3.8	C. 3	D. none of these

22. Absolute zero is the temperature at which

A. water solidifiesC. motion of molecules23. Motion of liquid in		http://www.upadmission.com/ B. all gases become liquid D. everything solidifies	
•	B. Poiseuille Equation (C. Stoke's Law	D. Archimedes' Principle
24. Molecular motion sA. Temperature25. Which is this gate ?A. ANDC. OR	B. Internal Energy	C. Friction	D. Viscosity
26. Energy bands in solA. Ohm's LawC. Bohr's Theory	lids are a consequence of	B. Pauli's Exclusion Pri D. Heissenberg's Uncer	-
a which is less than g. TA. Mg x Ma	The force exerted by the $B. g + a$	levator moving downwa boy on the floor of the el C. Mg – Ma er body B of mass m ₂ . If	levator is D. Mg + Ma
· ·	(in magnitude) of A is B. m_1m_2 a_2	C. m_1/m_2 (a ₂)	D. $(m_1 + m_2) a_2$
29. What does not chan A. Wavelength	ge when sound enters from B. Speed	om one medium to anoth C. Frequency	ner ? D. none of these
30. Resolving power of A. wavelength of light C. frequency of light us	<u>-</u>	npon B. wavelength of light of D. focal length of objections.	<u> </u>
31. An astronaut of wei apparent weight of the A. 5Kg	_	celerating upward with a	n acceleration of 4g. The D. zero
-	nters a magnetic field of at is the radius of the circ B. 100 m	10⁴ m/s normally, sp. chele describe by it ? C. 10 m	arge = 10 ¹¹ C/kg, D. none of these

33. If a black body radiates 20 calories per second at 227°C, it will radiate at 727°C

A. 10 calories per second	B. 80 calories per second	C. 320	http://www.upadmission.com/	
	per second	calories		
		per second D. none of these		
34. If a carnot engine is temperature is at 27°C,	s working with source te	mperature equal to 227°	C and its sink	
A. 20%	B. 10%	C. 67%	D. 50%	
35. If the frequency of energy is	an oscillating particle is	<i>n</i> , then the frequency of	oscillation of its potential	
A. n	B. 2n	C. n/2	D. 4n	
36. If an electron oscill A. X-rays C. Infra-red rays	lates at a frequency of 1	GHz, it gives : B. Micro-waves D. None of these		
37. Earth's atmosphere A. Ultra-violet rays	is richest in B. Infra-red rays	C. X-rays	D. Micro-waves	
38. Cathode rays consi	st of			
A. Photons	B. Electrons	C. Protons	D. α -particles	
<u> </u>	is moving with a velocity edded. At the point of co		ne system	
1 3	ring with velocity V in specomes stationary. What	, 0		
A. 4V	B. V	C. 4V/3	D. 2V/3	
41. A thief steals a box of weight W & jumps from the third floor of a building. During jump, he experiences a weight of				
A. W	B. 3W	C. 1.5W	D. zero	
42. Two electron beam A. they will attract each C. no interaction will to		space but in opposite dir B. they will repel each D. none of these		

43. Two wires with resistances R and 3R are connected in parallel, the ratio of heat generated in

2R and R is

A. 1:3 B. 2:1 C. 1:4

44. A wire is drawn such that its radius changes from r to 2r, the new resistance is

D. 4:1

A. 2 times	B. 4 times	http://www.u	npadmission.com/ D. 1/16 times		
45. In solids, inter-atom A. totally repulsive C. combination of (a) a		B. totally attractive D. none of these			
46. When horse starts running all of a sudden, the rider on the horse back falls backward because A. he is taken abackB. he is afraidC. due to inertia of rest, the upper part of his body remains at restD. due to inertia of motion, the lower part of his body comes in motion					
47. What should be the the string just does not	minimum velocity at the	highest point of a body	tied to a string, so that		
A. $\sqrt{(Rg)}$	B. $\sqrt{(5Rg)}$	C. $(R/g)^{3/2}$	D. $\sqrt{(2Rg)}$		
48. If a person standing A. increase C. remain same	on a rotating disc stretch	nes out his hands, the spe B. decrease D. none of these	eed will:		
49. EMF is most closely A. mechanical force	y related to B. potential difference (C. electric field	D. magnetic field		
50. Planetary system in A. conservation of ener C. conservation of angu		es B. conservation of linea D. none of these	ar momentum		
51. Lenz's law is based	upon				
A. energy	B. momentum	C. angular momentum	D. inertia		
52. Faraday's second la	w states that mass deposi B. atomic mass x	ted on the electrode is d	irectly proportional to		
A. atomic mass	velocity	C. atomic mass/valency	D. valency		
53. Unit of power is					
A. kilowatt hour	B. kilowatt per hour	C. kilowatt	D. erg		
54. Power can be expre A. F.v	ssed as B. 1/2 (Fv²)	55. Units of coefficient of viscosity are	C. F.t		

D. F x v

A. Nms⁻¹

B. Nm^2s^{-1}

 $C.\ Nm^{\text{--}2}\ s$

D. Nms⁻²

56. Dimensions of torq A. MLT ² B. ML ² T ²	ue are	C. M ² L ² T ² D. ML ⁻² T ²		
57. A body of weight <i>n</i> extending the string is	ag is hanging on a string,	which extends its length	by l . The work done in	
A. mg l	B. mg l/2	C. 2 mg l	D. none of these	
58. The water droplets	in free fall are spherical	due to		
A. gravity	B. viscosity	C. surface tension	D. inter-molecular attraction	
59. A ball of mass 1Kg A. 1 Kg ms ⁻²	is accelerating at a rate of B. 2 Kg ms ⁻²	of 1ms ⁻² . The rate of char C. 3 Kg ms ⁻²	nge of momentum is D. 4 Kg ms ⁻²	
60. A body orbitting are orbit of a satellite. The	ound earth at a mean rad period of the body is	ius which is two times as	s great as the parking	
A. 4 days	B. $2\sqrt{2}$ days	C. 16 days	D. 64 days	
61. Gamma rays are A. high energy electron C. high energy electro-		B. low energy electrons D. high energy positrons		
	abundant metal in the ear		5 W	
A. Fe	B. Al	C. Ca	D. Na	
	t give a precipitate with			
A. ZnSO ₄	B. FeSO ₄	C. AgNO ₃	D. HgCl ₂	
64. What volume of CO ₂ will be liberated at NTP of 12 gm of carbon is burnt in excess of oxygen?				
A. 11.2 litres	B. 22.4 litres	C. 2.24 litres	D. 1.12 litres	
65. Which base is found A. Adenine	d only in nucleotides of l B. Uracil	RNA? C. Guanine	D. Cytosine	
66. Ascorbic acid is the A. Vitamin B ₆	e chemical name of B. Vitamin A	C. Vitamin C	D. Vitamin D	

67. A hydrocarbon has carbon and hydrogen. Its molecular weight is 28. Its possible formula

would be

A. C ₃ H ₆	B. C ₂ H ₄	http://www.	upadmission.com/ D. C4H8		
68. The first Noble Priz	e in chemistry was giver B. Cnrizzaro	n to C. Mendeleevs	D. Moseley		
69. Four different collor action?	ids have the following go	old number. Which one	has its most effective		
A. 10	B. 30	C. 20	D. 40		
70. Which is an exampl	e of thermosetting polyn	ner?			
A. Polythene	B. PVC	C. Neoprene	D. Bakelite		
71. The number of unpa A. 3	nired electrons in ferrous B. 2	ion is C. 4	D. 5		
72. Strongest reducing a A. K	agent is B. Mg	C. Al	D. Ba		
73. Which of the follow	ring is man-made elemer	nt?			
A. Ra	B. U	C. Np	D. C – 4		
74. Which of the following statements is/are correct? A. Boiling point of alkylhalide is greater than its corresponding alkane B. In water, solubility of CH ₃ OH > C ₂ H ₅ OH > C ₆ H ₅ OH C. Aniline is a weaker base than NH ₃ D. All of the above					
75. Which amine of the A. Ethylamine	following will not answ B. Methylamine	er Carbylamine reaction C. Dimethylamine	? D. Phenylamine		
76. Tollen's reagent can A. (CH ₃) ₂ – CHOH	be used to detect B. CH ₃ – CO.CH ₃	C. CH ₃ CH ₂ CHO	D. CH ₃ OCH ₃		
77. Glycerol on heating A. Acetone	with Potassium bisulpha B. Glyceraldehyde	ate yields C. Acrolein	D. Propanol		
78. Salicylic acid on hea	ating with sodalime give B. Calcium salicylate		D. Phenol 79. Which one of the		

following will not give iodoform test?

A. Ethanol

B. Ethanal

C. 2-propanone

D. None of these

80. The rusting of iron A. Fe	is catalysed by B. O2	C. Zn	D. H ⁺
-		of a liquid B to give non	-ideal solution of A-B
mixture. The volume of A. 75 ml	tills illixture will be	B. 125 ml exact	
C. fluctuating between	75 ml and 125 ml	D. close to 125 ml but	not to exceed 125 ml
	•	mula $(CH_3)_3 C - CH = C$	
A. 3, 3 - dimethyl - 1 - b		B. 1, 1 - dimethyl - 3 -	
C. 1,1, 1 - dimethyl - 2	- propene	D. 3, 3, 3 - dimethyl - 1	- 1 propene
83. Which of the follow	ving compounds will be	optically active?	
A. $(OH_3)_2$ – CHOH	B. CH ₃ - CH ₂ - CH ₂ - CH ₃	C. CH ₃ – CHCl.COOH	D. (CH ₃) ₃ .C.Cl
84. The major compone	ents of brass are		
A. Zn and Sn	B. Cu and Zn	C. Fe and Ni	D. Zn and Fe
85. Lunar castic is			
A. Silver Chloride	B. Silver Nitrate	C. Sodium Hydroxide l	D. Potassium Nitrate
86. When hot iron is ex	posed in hot water vapo	ur, the compound formed	d is
A. FeO	B. Fe ₂ O ₄	C. Fe ₃ O ₄	D. Fe ₂ (OH) ₂
87. Which of the follow	ving halide is not oxidise	ed by MnO ₂ ?	
A. F	B. Cl C. Br	·	D. I
88. The outermost elect	tronic configuration of th	ne most electronegative e	element is
A. ns ² np ³	B. ns ² np ⁴	C. ns ² np ⁵	D. ns ² np ⁶
89. Shape of CO ₂ is			
A. tetrahedral	B. trigonal	C. bent	D. linear
90. The catalyst used in	the manufacture of H ₂ S	O ₄ by contact process is	
A. Al ₂ O ₃	B. Cr ₂ O ₃	C. V ₂ O ₅	D. MnO ₂
91. The composition of	the common glass is		
A. Na ₂ O.CaO.6SiO ₂ B. 1	Na ₂ O.Al ₂ O ₃ .2SiO ₂ C. Ca	$10.Al_2O_3.2SiO_2D.$	Na ₂ O.CaO.Al ₂ O ₃ .6SiO ₂

92. In a borax lead test, the brown colour is due to http://www.upadmission.com/				
A. Chromium	B.Cobalt	C. Manganese	D. Iron	
93. Which of the follow A. Urea	ving is not a fertiliser? B. Superphosphate of lime	C. Benzene Hexachloride	D. Potassium	
94. Which one of the formula Table?	ollowing belongs to repr	esentative group of elem	ents in the Periodic	
A. Lanthanum	B. Argon	C. Chromium	D. Aluminium	
95. Which one of the fo	ollowing is not an isotop B. Deuterium	e of Hydrogen? C. Ortho-hydrogen	D. None of the above	
96. In the reaction $I_2 + 2S_2O_{32} = 2I^2 + S_4O_6^{22}$, equivalent weight of iodine will be equal to A. its molecular weight B. 1/2 of its molecular weight C. 1/4 the molecular weight D. twice the molecular weight				
97. Which of the follow A. F ₂ C. Br ₂	ving is the most powerfu	l oxidising agent? B. Cl ₂ D. I ₂		
98. From the following strongest acid?	values of dissociating c	onstants of four acids, w	hich value represents the	
A. 2×10^{-2}	B. 0.02 x 10 ⁻¹	C. 3 x 10 ⁻³	D. 2.0 x 10 ⁴	
99. In which of the foll A. $K = 10^3$	owing cases, does the re B. $K = 10^{-2}$	action go the farthest for C. K = 10	completion? D. K = 1	
100. The reaction which proceeds in the forward direction is A. Fe ₂ O ₃ + 6HCl \rightarrow 2FeCl ₃ + 3H ₂ O B. NH ₃ + H ₂ O + NaCl \rightarrow NH ₄ Cl + NaOH C. SnCl ₄ + Hg ₂ Cl ₂ \rightarrow SnCl ₂ + 2HgCl ₂ D. 2CuI + I ₂ + 4K ⁺ \rightarrow 2Cu ²⁺ + 4KI				
101. The substance cap A. malleable	bable of being drawn into B. tensile	o fine wire is called C. ductile	D. mild	
102. The idea that mos is given by	t of the mass of an atom	is concentrated in a very	small core, i.e., nucleus	
A. Amedo Avogadro	B. Rutherford	C. Bohr	D. Henery Mosley	
103. Which of the follo	owing does contain a co-	ordinate covalent bond?	A. N ₂ H ₅₊ BaCl ₂ C. HCl	

B.

	,		. ,
http://	/www.upa	dmissi	ion.com/

		-	upadmission.com/
	owing contains both cova		
A. CCl ₄	B. CaCl ₂	C. NH ₄ Cl D. H ₂ O	
	he periodic law and the phe maximum electronega	periodic table, suggest wlative character?	hich of the following
A. Oxygen	B. Nitrogen	C. Fluorine	D. Astatine
106. The electronic cor	nfiguration of element ato	omic number 37 is	
A. $(2, 8) 3s^23p^63d^{10}4s^24p^2$	p^65s^1	B. $(2, 8) 3s^23p^63d^{10}4s^25s$	$^{6}4p^{5}$
C. $(2, 8) 3s^23p^64s^23d^95s^1$	¹ 4p ⁵	D. none of these	
107. The pH of 0.1 M s the acid?	solution of a weak acid is	s 3. What is the value of	ionisation constant for
A. 0.1	B. 10 ⁻³	C. 10 ⁻⁵	D. 10 ⁻⁷
108. Pure Aniline is a			
A. brown coloured liquid	B. colourless liquid	C. brown coloured solid	D. colourless solid
109. Sulphide ores are A. roasting	generally concentrated b B. froth floatation	y C. reducing by carbon l	D. tempering
110. One mole of CO ₂	contains		
A. 6.02×10^{23} atoms of		B. 6.02×10^{23} atoms of	O
C. 18.1 x 10 ²³ molecule	es of CO ₂	D. 3 gm atom of CO ₂	
111. The Avogadro Nu	umber or a mole represen	ts 23	
A. 6.02 x 10 ²³ ions	B. 6.02 x 10 ²³ atoms	C. 6.02 x 10 molecules	D. 6.02 x 10 ²³ entities
112. What is the weigh	at of one molecule of a m	onoatomic element X wh	nose atomic weight is 36?
A. 6.0 x 10 ⁻²³ gm	B. 6.02 x 10 ²³ gm	C. 36 x 10 ²³ gm	D. 36 x 10 ⁻²³ gm
113. When α -particles because	are set through a thin me	etal foil, most of them go	straight through the foil
A. α -particles are muc	h heavier than electrons	B. α -particles are posit	ively charged
C. α -particles move with	ith high velocity	D. α -particles move with	ith low velocity
114. The reaction, which	ch proceeds in the forwar	rd direction, is	
A. $Fe_2O_3 + 6HCl \rightarrow 2F$	$FeCl_3 + 3H_2O$	C. SnCl ₄ + Hg ₂	$Cl_2 \rightarrow SnCl_2 + 2HgCl_2$

http://www.upadmission.com/

115. The first order coperiod for this decom	onstant for the decomposition in second is	ition of N_2O_5 is 6.2×10^4	sec –1. The half-life			
A. 1117.7	B. 111.7	C. 223.4	D. 160.9			
116. When the same amount of zinc is treated separately with excess of H2SO4 and excess of						
NaOH, the ratio of vo A. 1:1	blumes of H ₂ evolved is B. 1:2	C. 2:1	D. 9:4			
11.1.1	2.1.2	0.2.1	2.7.1			
	ot combine directly with	0.1.1	D 1			
A. oxygen	B. nitrogen	C. hydrogen	D. carbon			
118. Carbon differs fr	com other elements of its	sub-group due to				
A. availability of d-or	· ·	B. its limitation to a co-ordination number four				
C. its tendency to cate	enate	D. its unique ability to form multiple bonds				
119. Iodine reacts wit	h cold dil. NaOH to give					
A. $NaI + H_2O + O_2$	B. $NaI + NaIO + O_2C$.	$NaI + NaIO + H_2O D. N$	$VaI + NaIO_3 + H_2O$			
120. The number of is	somers for the atomic cor	npound of the formula C	7H8O is			
A. 2	B. 3	C. 4	D. 5			
121. Which of the fol	lowing is not true in linea	ar programming problem	?			
A. A column in the						
simplex table that						
contains all of the						
variables in the soluti is called pivot or key	on					
column.						
B. A basic solution						
which is also in the						
feasible region is call-	ed					
a basic feasible						
solution.						
C. A surplus variable	is					
a variable subtracted						
from the left hand sid	e					
of a greater than or						
equal to constraint to convert it into an						
equality.						

D. A slack variable is a

variable added to the




constraint to convert i into an equality.	t				
122. The equation of topasses through (4, 6) is		er lies on $2x + 3y = 3$ and	116x - y = 4 and which		
A. $x^2 + y^2 = 40$		B. $5(x^2 + y^2) - 4x - 8y$	B. $5(x^2 + y^2) - 4x - 8y = 200$		
C. $x^2 + y^2 - 4x - 8y = 200$		· • •	D. $5(x^2 + y^2) - 3x - 8y = 200$		
123. Let $n(A) = 4$ and A. 120	n(B) = 5. The number of B. 9	of all possible injections of C. 24	from A to B is D. none		
124. If $aN = \{ax : x \in$	N } and $bN \cap cN = dN$, where b, $c \in N$ are relati	ively prime, then		
A. $c = bd$	B. b = cd	C. d = bc	D. none of the above		
125. A square root of	3 + 4i is				
A. $\sqrt{3} + i$	B. 2 - i	C. $2 + i$	D. none of the above		
126. Which of the foll A.	lowing is not applicable C. D.	for a complex number?			
Inequality B. Division	on Subtraction Addition				
127. maximum amp	(z) - minimum amp (z)	is equal to			
A. sin ⁻¹ (3/5) - cos ⁻¹ (3/	(5)	B. $\pi/2 + \cos^{-1}(3/5)$			
C. π - 2 $\cos^{-1}(3/5)$		D. $\cos^{-1}(3/5)$			
128. If e, e' be the ecc be	entricities of two conics	s S and S' and if $e^2 + e^{t^2} =$	3, then both S and S' can		
A. hyperbolas	B. ellipses	C. parabolas	D. none of the above		
=	'l' rests against the floo us of its middle point is	r and a wall of a room. If	the stick begins to slide on		
A. an ellipse	B. a parabola	C. a circle	D. a straight line		
= = = = = = = = = = = = = = = = = = = =	-	ets the straight line x/y + of y and whose axes lie alo			
A. $2\sqrt{6/7}$	B. $3\sqrt{2/7}$	C. √6/7	D. none of the above		
-	tive acute angles satisfy A, then A + 2B is equa	<u> </u>	$A + 2\cos^2 B = 4 \text{ and } 3 \sin \theta$		

C. $\pi/6$

D. $\pi/4$

Β. π/2

A. $\pi/3$



B.

C.

D. none of the above

http://www.upadmission.com/

144. How many different arrangements can be made out of the letters in the expansion A²B³C⁴, when written in full?

145. The number of straight lines that can be drawn out of 10 points of which 7 are collinear is

146. $1/n! + 1/[2! (n-2)!] + 1/[4! (n-4)!] + \dots$ is

A.
$$(2n-1)/n!$$

B.
$$2^{n}/[(n+1)!]$$

B.
$$2^{n}/[(n+1)!]$$
 C. $2^{n}/n!$ D. $2^{n-2}/[(n-1)!]$

147. The term independent of x in $(x^2 - 1/x)^9$ is

D. none of the above

148. The 9th term of an A.P. is 499 and 499th term is 9. The term which is equal to zero is

A. 501th

D. none of the above

A. is a skew symmetric matrix

$$B. A^{-1} + B^{-1}$$

C. does not exist

D. none of the above


150. If AB = A and BA = B, then B^2 is equal to

A. B

B. A

C. 1

D. 0



A. a, b, c are in H.P.

B. α is a root of $4ax^2 + 12bx + 9c = 0$ or a, b, c are in G.P.

C. a, b, c are in G.P. only

a, b, c are in A.P.

http://www.upadmission.com/

152. The value of K so that (x - 1)/-3 = (y - 2)/2K = (z - 3)/2 and (x - 1)/3K = (y - 1)/1 = (z - 6)/-15 may be perpendicular is given by

- A. -7/10
- B. -10/7
- C. -10

D. 10/7

153. The equation of the plane containing the line

A.
$$+\mathbf{i} \cdot (+\mathbf{i} - 2\mathbf{j} + \mathbf{k}) = 0$$

B.
$$(\stackrel{i}{\rightarrow} 2\stackrel{i}{\rightarrow} -k) \rightarrow 0 \rightarrow$$

C.
$$(i + 2j - k) = 3$$

D. none of the above

154. The mean of discrete observations y_1, y_2, \dots, y_n is given by $\sum_{i=1}^{n} y_i f_i$

$$\sum_{i=1}^{n} y_i f_i$$

 $y_{\scriptscriptstyle i}\;f_{\scriptscriptstyle i}$

A.
$$\sum_{i=1}^{n} \frac{f_i}{n}$$

B.
$$\sum_{i=1}^{n} n$$

Уi

 $y_{\rm i}$


C.

B.

i

n

155. For a poisson	distribution	whose mean is λ , the standard	deviation will be
A. λ^2	Β. 1/λ	C. $\sqrt{\lambda}$	D. λ



A. 4	B. 1	C. 9	D. 8		
168. The area bounded A. 32/7	by the curve $y^2 = 8x$ and B. 24/5	$x^2 = 8y$ is C. $72/3$	D. 64/3		
169. The integrating factory	ctor of the differential eq	uation $[(dy/dx)(x \log x)]$	$+ y = 2 \log x$ is given		
A. $\log(\log x)$	B. e ^x	C. log x	D. x		
170. If $y = tan^{-1}[(\sin x + A. 1/2)]$	$\cos x$ /($\cos x - \sin x$)], the B. 0	nen dy/dx is equal to C. 1	D. none of the above		
171. The length of tang A. 81	ent from (5, 1) to the circ B. 29	cle $x^2 + y^2 + 6x - 4y - 3 =$ C. 7	0 is D. 21		
172. The equation of the straight line which is perpendicular to $y = x$ and passes through $(3, 2)$					
will be given by A. $x - y = 5$	B. $x + y = 5$	C. $x + y = 1$	D. $x - y = 1$		
173. If the imaginary part of $(2z + 1)/(iz + 1)$ is - 2, then the locus of the point representing z in the complex plane is					
A. a circle	B. a straight line	C. a parabola	D. none of the above		
174. The sum of 40 term A. 3200	ns of an A.P. whose first B. 1600	term is 2 and common of C. 200	lifference 4, will be D. 2800		
175. If a, b, c are in A.F. A. A.P.	P., then a/bc, 1/c, 2/b are B. G.P.	in C. H.P.	D. none of the above		
176. The term independent of x in $[x^2 + (1/x^2)]$ is					
A. 1	B1	C. 48	D. none of the above		
177. The equation of a A . $y = -3$	line through $(2, -3)$ paral B. $y = 2$	lel to y-axis is C. $x = 2$	D. $x = -3$		
178. The value of (ax)	$^{3} + bx + c$) dx depends				
A. the value of b	B. the value of c	C. the value of a	D. the value of a and b		

167. The least value of 'a' for which the equation $(4/\sin x) + [1/(1 - \sin x)] = a$ has at least one

solution on the interval $(0, \pi/2)$ is

179. The range of the function $f(x) = (1 + x^2)/x^2$ is equal to **http://www.upadmission.com/** B. [1, 0] $C.(1,\infty)$ A. [0, 1] D. $[2, \infty]$

180. Two vectors are said to be equal if

B. direction is same A. their magnitudes are same

D. they have magnitude and same sense of C. they meet at the same point

direction