

## WBJEE (Engineering) 2015


### Solution

#### PHYSICS

##### Category – I (Q1 to Q30)

Each question has one correct option and carries 1 mark, for each wrong answer 1/4 mark will be deducted.

1. An object is located 4 m from the first of two thin converging lenses of focal lengths 2 m and 1 m respectively. The lenses are separated by 3 m. The final image formed by the second lens is located from the source at a distance of
- (A) 8.0 m                      (B) 7.5 m                      (C) 6.0 m                      (D) 6.5 m



**Solution : (B)**

2. A simple pendulum of length  $L$  swings in a vertical plane. The tension of the string when it makes an angle  $\theta$  with the vertical and the bob of mass  $m$  moves with a speed  $v$  is ( $g$  is the gravitational acceleration)
- (A)  $mv^2/L$                       (B)  $mg \cos\theta + mv^2/L$                       (C)  $mg \cos\theta - mv^2/L$                       (D)  $mg \cos\theta$

**Solution : (B)**

3. The length of a metal wire is  $L_1$  when the tension is  $T_1$  and  $L_2$  when the tension is  $T_2$ . The unstretched length of the wire is

(A)  $\frac{L_1 + L_2}{2}$                       (B)  $\sqrt{L_1 L_2}$                       (C)  $\frac{T_2 L_1 - T_1 L_2}{T_2 - T_1}$                       (D)  $\frac{T_2 L_1 + T_1 L_2}{T_2 + T_1}$

**Solution : (C)**

4. A hollow sphere of external radius  $R$  and thickness  $t$  ( $t \ll R$ ) is made of a metal of density  $\rho$ . The sphere will float in water if

(A)  $t \leq \frac{R}{\rho}$                       (B)  $t \leq \frac{R}{3\rho}$                       (C)  $t \leq \frac{R}{2\rho}$                       (D)  $t \geq \frac{R}{3\rho}$


**Solution : (B)**

5. A metal wire of circular cross-section has a resistance  $R_1$ . The wire is now stretched without breaking so that its length is doubled and the density is assumed to remain the same. If the resistance of the wire now becomes  $R_2$  then  $R_2 : R_1$  is

(A) 1 : 1                      (B) 1 : 2                      (C) 4 : 1                      (D) 1 : 4

**Solution : (C)**

6. Assume that each diode shown in the figure has a forward bias resistance of  $50\Omega$  and an infinite reverse bias resistance. The current through the resistance  $150\Omega$  is  
 (A) 0.66 A (B) 0.05 A (C) Zero (D) 0.04 A



**Solution : (D)**


7. A straight conductor 0.1m long moves in a uniform magnetic field 0.1T. The velocity of the conductor is 15 m/s and is directed perpendicular to the field. The e.m.f. induced between the two ends of the conductor is  
 (A) 0.10 V (B) 0.15 V (C) 1.50 V (D) 15.00 V

**Solution : (B)**

8. A ray of light is incident at an angle  $i$  on a glass slab of refractive index  $\mu$ . The angle between reflected and refracted light is  $90^\circ$ . The relationship between  $i$  and  $\mu$  is  
 (A)  $i = \tan^{-1}\left(\frac{1}{\mu}\right)$  (B)  $\tan i = \mu$  (C)  $\sin i = \mu$  (D)  $\cos i = \mu$


**Solution : (B)**

9. Two particles A and B are moving as shown in the figure. Their total angular momentum about the point O is  
 (A)  $9.8 \text{ kg m}^2/\text{s}$  (B) Zero (C)  $52.7 \text{ kg m}^2/\text{s}$  (D)  $37.9 \text{ kg m}^2/\text{s}$



**Solution : (A)**

10. Particle A moves along X-axis with a uniform velocity of magnitude 10 m/s. Particle B moves with uniform velocity 20 m/s along a direction making an angle of  $60^\circ$  with the positive direction of X-axis as shown in the figure. The relative velocity of B with respect to that of A is  
 (A) 10 m/s along X-axis  
 (B)  $10\sqrt{3}$  m/s along Y-axis (Perpendicular to X-axis)  
 (C)  $10\sqrt{5}$  m/s along the bisection of the velocities of A and B  
 (D) 30 m/s along negative X-axis




**Solution : (B)**

11. When light is refracted from a surface, which of its following physical parameters does not change ?  
 (A) velocity (B) amplitude (C) frequency (D) wavelength  
**Solution : (C)**
12. A solid maintained at  $t_1^\circ \text{C}$  is kept in an evacuated chamber at temperature  $t_2^\circ \text{C}$  ( $t_2 \gg t_1$ ). The rate of heat absorbed by the body is proportional to  
 (A)  $t_2^4 - t_1^4$  (B)  $(t_2^4 + 273) - (t_1^4 + 273)$   
 (C)  $t_2 - t_1$  (D)  $t_2^2 - t_1^2$   
**Solution : (C)**
13. The work function of metals is in the range of 2 eV to 5eV. Find which of the following wavelength of light cannot be used for photoelectric effect. (Consider, Plank constant =  $4 \times 10^{-15}$  eVs, velocity of light =  $3 \times 10^8$  m/s)  
 (A) 510 nm (B) 650 nm (C) 400 nm (D) 570 nm  
**Solution : (B)**
14. A thin plastic sheet of refractive index 1.6 is used to cover one of the slits of a double slit arrangement. The central point on the screen is now occupied by what would have been the 7<sup>th</sup> bright fringe before the plastic was used. If the wavelength of light is 600 nm, what is the thickness (in  $\mu\text{m}$ ) of the plastic?  
 (A) 7 (B) 4 (C) 8 (D) 6  
**Solution : (A)**
15. The length of an open organ pipe is twice the length of another closed organ pipe. The fundamental frequency of the open pipe is 100 Hz. The frequency of the third harmonic of the closed pipe is  
 (A) 100 Hz (B) 200 Hz (C) 300 Hz (D) 150 Hz  
**Solution : (C)**
16. A  $5 \mu\text{F}$  capacitor is connected in series with a  $10 \mu\text{F}$  capacitor. When a 300 Volt potential difference is applied across this combination, the total energy stored in the capacitors is  
 (A) 15 J (B) 1.5 J (C) 0.15 J (D) 0.10 J  
**Solution : (C)**
17. Two particles of mass  $m_1$  and  $m_2$ , approach each other due to their mutual gravitational attraction only. Then  
 (A) accelerations of both the particles are equal.  
 (B) acceleration of the particle of mass  $m_1$  is proportional to  $m_1$ .  
 (C) acceleration of the particle of mass  $m_1$  is proportional to  $m_2$ .  
 (D) acceleration of the particle of mass  $m_1$  is inversely proportional to  $m_1$ .  
**Solution : (C)**
18. Three bodies of the same material and having masses  $m$ ,  $m$  and  $3m$  are at temperature  $40^\circ\text{C}$ ,  $50^\circ\text{C}$  and  $60^\circ\text{C}$  respectively. If the bodies are brought in thermal contact, the final temperature will be  
 (A)  $45^\circ\text{C}$  (B)  $54^\circ\text{C}$  (C)  $52^\circ\text{C}$  (D)  $48^\circ\text{C}$   
**Solution : (B)**
19. A satellite has kinetic energy  $K$ , potential energy  $V$  and total energy  $E$ . Which of the following statements is true?  
 (A)  $K = -V/2$  (B)  $K = V/2$  (C)  $E = K/2$  (D)  $E = -K/2$   
**Solution : (A)**

20. The line AA' is on a charged infinite conducting plane which is perpendicular to the plane of the paper. The plane has a surface density of charge  $\sigma$  and B is a ball of mass  $m$  with a like charge of magnitude  $q$ . B is connected by a string from a point on the line AA'. The tangent of the angle ( $\theta$ ) formed between the line AA' and the string is :


- (A)  $\frac{q\sigma}{2\epsilon_0 mg}$       (B)  $\frac{q\sigma}{4\pi\epsilon_0 mg}$       (C)  $\frac{q\sigma}{2\pi\epsilon_0 mg}$       (D)  $\frac{q\sigma}{\epsilon_0 mg}$



**Solution : (D)**

21. The current  $I$  in the circuit shown is

- (A) 1.33 A      (B) Zero      (C) 2.00 A      (D) 1.00 A



**Solution : (A)**

22. The r.m.s speed of oxygen is  $v$  at a particular temperature. If the temperature is doubled and oxygen molecules dissociate into oxygen atoms, the r.m.s speed becomes

- (A)  $v$       (B)  $\sqrt{2}v$       (C)  $2v$       (D)  $4v$

**Solution : (C)**

23. Two particles, A and B, having equal charges, after being accelerated through the same potential difference enter a region of uniform magnetic field and the particles describe circular paths of radii  $R_1$  and  $R_2$  respectively. The ratio of the masses of A and B is

- (A)  $\sqrt{R_1/R_2}$       (B)  $R_1/R_2$       (C)  $(R_1/R_2)^2$       (D)  $(R_2/R_1)^2$

**Solution : (C)**

24. A large number of particles are placed around the origin, each at a distance  $R$  from the origin. The distance of the centre of mass of the system from the origin is

- (A)  $= R$       (B)  $\leq R$       (C)  $> R$       (D)  $\geq R$

**Solution : (B)**

25. A 20 cm long capillary tube is dipped vertically in water and the liquid rises upto 10 cm. If the entire system is kept in a freely falling platform, the length of water column in the tube will be

- (A) 5 cm      (B) 10 cm      (C) 15 cm      (D) 20 cm

**Solution : (D)**

26. A train is moving with a uniform speed of 33 m/s and an observer is approaching the train with the same speed. If the train blows a whistle of frequency 1000 Hz and the velocity of sound is 333 m/s, then the apparent frequency of the sound that the observer hears is

(A) 1220 Hz                      (B) 1099 Hz                      (C) 1110 Hz                      (D) 1200 Hz

**Solution : (A)**


27. A photon of wavelength 300 nm interacts with a stationary hydrogen atom in ground state. During the interaction, whole energy of the photon is transferred to the electron of the atom. State which possibility is correct. (Consider, Plank constant =  $4 \times 10^{-15}$  eVs, velocity of light =  $3 \times 10^8$  m/s, ionization energy of hydrogen = 13.6 eV)

(A) Electron will be knocked out of the atom  
 (B) Electron will go to any excited state of the atom  
 (C) Electron will go only to first excited state of the atom  
 (D) Electron will keep orbiting in the ground state of atom

**Solution : (D)**

28. Block B lying on a table weighs  $W$ . The coefficient of static friction between the block and the table is  $\mu$ . Assume that the cord between B and the knot is horizontal. The maximum weight of the block A for which the system will be stationary is


(A)  $\frac{W \tan \theta}{\mu}$                       (B)  $\mu W \tan \theta$                       (C)  $\mu W \sqrt{1 + \tan^2 \theta}$                       (D)  $\mu W \sin \theta$



**Solution : (B)**

29. The inputs to the digital circuit are shown below. The output Y is

(A)  $A + B + \bar{C}$                       (B)  $(A + B)\bar{C}$                       (C)  $\bar{A} + \bar{B} + \bar{C}$                       (D)  $\bar{A} + \bar{B} + C$



**Solution : (C)**


30. Two particles A and B having different masses are projected from a tower with same speed. A is projected vertically upward and B vertically downward. On reaching the ground
- (A) velocity of A is greater than that of B  
 (B) velocity of B is greater than that of A  
 (C) both A and B attain the same velocity  
 (D) the particle with the larger mass attains higher velocity

**Solution : (C)**

**Category – II (Q31 to Q35)**

Each question has one correct option and carries 2 marks, for each wrong answer 1/2 mark will be deducted.

31. Two cells A and B of e.m.f. 2V and 1.5V respectively, are connected as shown in figure through an external resistance  $10\Omega$ . The internal resistance of each cell is  $5\Omega$ . The potential difference  $E_A$  and  $E_B$  across the terminals of the cells A and B respectively are
- (A)  $E_A = 2.0V, E_B = 1.5V$  (B)  $E_A = 2.125V, E_B = 1.375V$   
 (C)  $E_A = 1.875V, E_B = 1.625V$  (D)  $E_A = 1.875V, E_B = 1.375V$



**Solution : (C)**

32. A charge  $q$  is placed at one corner of a cube. The electric flux through any of the three faces adjacent to the charge is zero. The flux through any one of the other three faces is
- (A)  $q/3\epsilon_0$  (B)  $q/6\epsilon_0$  (C)  $q/12\epsilon_0$  (D)  $q/24\epsilon_0$
33. In the circuit shown below, the switch is kept in position 'a' for a long time and is then thrown to position 'b'. The amplitude of the resulting oscillating current is given by
- (A)  $E\sqrt{L/C}$  (B)  $E/R$  (C) Infinity (D)  $E\sqrt{C/L}$

**Solution : (D)**


34. The pressure  $p$ , volume  $V$  and temperature  $T$  for a certain gas are related by  $p = \frac{AT - BT^2}{V}$ , where  $A$  and  $B$  are constants. The work done by the gas when the temperature changes from  $T_1$  to  $T_2$  while the pressure remains constant, is given by

- (A)  $A(T_2 - T_1) + B(T_2^2 - T_1^2)$  (B)  $\frac{A(T_2 - T_1)}{V_2 - V_1} - \frac{B(T_2^2 - T_1^2)}{V_2 - V_1}$   
 (C)  $A(T_2 - T_1) - B(T_2^2 - T_1^2)$  (D)  $\frac{A(T_2 - T_1^2)}{V_2 - V_1}$

**Solution : (C)**



35. A cylinder of height  $h$  is filled with water and is kept on a block of height  $h/2$ . The level of water in the cylinder is kept constant. Four holes numbered 1, 2, 3 and 4 are at the side of the cylinder and at heights  $0$ ,  $h/4$ ,  $h/2$  and  $3h/4$  respectively. When all four holes are opened together, the hole from which water will reach farthest distance on the plane  $PQ$  is the hole no.
- (A) 1                      (B) 2                      (C) 3                      (D) 4



**Solution :** (B)

**Category – III (Q36 to Q40)**

Each question has one or more correct option(s) choosing which will fetch maximum 2 marks on pro rata basis. However, choice of any wrong option(s) will fetch zero mark for the question.

36. Consider two particles of different masses. In which of the following situations the heavier of the two particles will have smaller de Broglie wavelength ?
- (A) Both have a free fall through the same height.    (B) Both move with the same kinetic energy.  
(C) Both move with the same linear momentum.    (D) Both move with the same speed.

**Solution :** (A, B, D)

37. A circular disc rolls on a horizontal floor without slipping and the centre of the disc moves with a uniform velocity  $v$ . Which of the following values the velocity at a point on the rim of the disc can have ?
- (A)  $v$                       (B)  $-v$                       (C)  $2v$                       (D) Zero

**Solution :** (A, C, D)

38. A conducting loop in the form of a circle is placed in a uniform magnetic field with its plane perpendicular to the direction of the field. An e.m.f. will be induced in the loop if
- (A) it is translated parallel to itself.  
(B) it is rotated about one of its diameters.  
(C) it is rotated about its own axis which is parallel to the field.  
(D) the loop is deformed from the original shape.

**Solution :** (B, D)

39. Find the right condition(s) for Fraunhofer diffraction due to a single slit.
- (A) Source is at infinite distance and the incident beam has converged at the slit.  
(B) Source is near to the slit and the incident beam is parallel.  
(C) Source is infinity and the incident beam is parallel.  
(D) Source is near to the slit and the incident beam has covered at the slit.

**Solution :** (B, C)

40. Two charges  $+q$  and  $-q$  are placed at a distance 'a' in a uniform electric field. The dipole moment of the combination is  $2qa(\cos\theta \hat{i} + \sin\theta \hat{j})$ , where  $\theta$  is the angle between the direction of the field and the line joining the two charges. Which of the following statement(s) is/are correct ?
- (A) The torque exerted by the field on the dipole vanishes.
  - (B) The net force on the dipole vanishes.
  - (C) The torque is independent of the choice of coordinates.
  - (D) The net force is independent of 'a'.

**Solution :** (B, C, D)




**PATHFINDER**<sup>®</sup>  
Where Aspirations Meet Success







**CHEMISTRY**

**Category – I (Q41 to Q70)**


Each question has one correct option and carries 1 mark, for each wrong answer 1/4 mark will be deducted.







The major product of the above reaction is



**Solution : (B)**



The product of the above reaction is



**Solution : (C)**

43. For the reaction  $A + 2B \rightarrow C$ , the reaction rate is doubled if the concentration of A is doubled. The rate is increased by four times when concentrations of both A and B are increased by four times. The order of the reactions is

(A) 3

(B) 0

(C) 1

(D) 2

**Solution : (C)**

44. At a certain temperature, the value of the slope of the plot of osmotic pressure ( $\pi$ ) against concentration ( $C$  in  $\text{mol L}^{-1}$ ) of a certain polymer solution is  $291R$ . The temperature at which osmotic pressure is measured is ( $R$  is gas constant)

(A)  $271^\circ\text{C}$  (B)  $18^\circ\text{C}$  (C)  $564\text{ K}$  (D)  $18\text{ K}$

**Solution : (B)**

45. The rms velocity of  $\text{CO}$  gas molecules at  $27^\circ\text{C}$  is approximately  $1000\text{ m/s}$ . For  $\text{N}_2$  molecules at  $600\text{ K}$  the rms velocity is approximately

(A)  $2000\text{ m/s}$  (B)  $1414\text{ m/s}$  (C)  $1000\text{ m/s}$  (D)  $1500\text{ m/s}$

**Solution : (B)**

46. A gas can be liquefied at temperature  $T$  and pressure  $P$  provided

(A)  $T = T_C$  and  $P < P_C$  (B)  $T < T_C$  and  $P > P_C$  (C)  $T > T_C$  and  $P > P_C$  (D)  $T > T_C$  and  $P < P_C$

**Solution : (B)**

47. In a mixture, two enantiomers are found to be present in  $85\%$  and  $15\%$  respectively. The enantiomeric excess ( $e, e$ ) is

(A)  $85\%$  (B)  $15\%$  (C)  $70\%$  (D)  $60\%$

**Solution : (C)**

48. 1,4-dimethylbenzene on heating with anhydrous  $\text{AlCl}_3$  and  $\text{HCl}$  produces

(A) 1,2-dimethylbenzene (B) 1,3-dimethylbenzene  
(C) 1,2,3-trimethylbenzene (D) Ethylbenzene

**Solution : (A)**

49. 

The product of the above reaction is

(A)  (B)   
(C)  (D) 

**Solution : (C)**

50. Suppose the mass of a single  $\text{Ag}$  atom is ' $m$ '.  $\text{Ag}$  metal crystallizes in fcc lattice with unit cell of length ' $a$ '. The density of  $\text{Ag}$  metal in terms of ' $a$ ' and ' $m$ ' is

(A)  $\frac{4m}{a^3}$  (B)  $\frac{2m}{a^3}$  (C)  $\frac{m}{a^3}$  (D)  $\frac{m}{4a^3}$

**Solution : (A)**

51. For the reaction  $2\text{SO}_2(\text{g}) + \text{O}_2(\text{g}) \rightleftharpoons 2\text{SO}_3(\text{g})$  at  $300\text{K}$ , the value of  $\Delta G^\circ$  is  $-690.9\text{R}$ . The equilibrium constant value for the reaction at that temperature is ( $R$  is gas constant)

(A)  $10\text{ atm}^{-1}$  (B)  $10\text{ atm}$  (C)  $10$  (D)  $1$

**Solution : (A)**

52. At a particular temperature the ratio of equivalent conductance to specific conductance of a 0.01 (N) NaCl solution is

- (A)  $10^5 \text{ cm}^3$                       (B)  $10^3 \text{ cm}^3$                       (C)  $10 \text{ cm}^3$                       (D)  $10^5 \text{ cm}^2$

**Solution : (A)**

53. The units of surface tension and viscosity of liquids are respectively

- (A)  $\text{kg m}^{-1}\text{s}^{-1}$ ,  $\text{N m}^{-1}$     (B)  $\text{kg s}^{-2}$ ,  $\text{kg m}^{-1}\text{s}^{-1}$     (C)  $\text{N m}^{-1}$ ,  $\text{kg m}^{-1}\text{s}^{-2}$     (D)  $\text{kg s}^{-1}$ ,  $\text{kg m}^{-2}\text{s}^{-1}$

**Solution : (C)**

54. The ratio of volumes of  $\text{CH}_3\text{COOH}$  0.1 (N) to  $\text{CH}_3\text{COONa}$  0.1 (N) required to prepare a buffer solution of pH 5.74 is (given :  $\text{pK}_a$  of  $\text{CH}_3\text{COOH}$  is 4.74)

- (A) 10 : 1                      (B) 5 : 1                      (C) 1 : 5                      (D) 1 : 10

**Solution : (A)**

55. The reaction of methyltrichloroacetate ( $\text{Cl}_3\text{CCO}_2\text{Me}$ ) with sodium methoxide ( $\text{NaOMe}$ ) generates

- (A) Carbocation                      (B) Carbene                      (C) Carbanion                      (D) Carbon radical

**Solution : (B)**

56. Best reagent for nuclear iodination of aromatic compounds is

- (A)  $\text{KI}/\text{CH}_3\text{COCH}_3$                       (B)  $\text{I}_2/\text{CH}_3\text{CN}$                       (C)  $\text{KI}/\text{CH}_3\text{COOH}$                       (D)  $\text{I}_2/\text{HNO}_3$


**Solution : (B)**

57. In the Lassaigne's test for the detection of nitrogen in an organic compound, the appearance of blue coloured compound is due to

- (A) ferric ferricyanide                      (B) ferrous ferricyanide  
(C) ferric ferrocyanide                      (D) ferrous ferrocyanide

**Solution : (C)**

58. In the following reaction



The product 'P' is

- (A)  $\text{RCHO}$                       (B)  $\text{R}_2\text{CHOEt}$                       (C)  $\text{R}_3\text{CH}$                       (D)  $\text{RCH(OEt)}_2$

**Solution : (A)**

59. The dispersed phase and dispersion medium of fog respectively are

- (A) solid, liquid                      (B) liquid, liquid                      (C) liquid, gas                      (D) gas, liquid

**Solution : (C)**

60. The decreasing order of basic character of  $\text{K}_2\text{O}$ ,  $\text{BaO}$ ,  $\text{CaO}$  and  $\text{MgO}$  is

- (A)  $\text{K}_2\text{O} > \text{BaO} > \text{CaO} > \text{MgO}$                       (B)  $\text{K}_2\text{O} > \text{CaO} > \text{BaO} > \text{MgO}$   
(C)  $\text{MgO} > \text{BaO} > \text{CaO} > \text{K}_2\text{O}$                       (D)  $\text{MgO} > \text{CaO} > \text{BaO} > \text{K}_2\text{O}$

**Solution : (A)**

61. In aqueous alkaline solution, two electron reduction of  $\text{HO}_2^-$  gives

- (A)  $\text{HO}^-$                       (B)  $\text{H}_2\text{O}$                       (C)  $\text{O}_2$                       (D)  $\text{O}_2^-$

**Solution : (A)**

62. Cold ferrous sulphate solution on absorption of NO develops brown colour due to the formation of  
 (A) paramagnetic  $[\text{Fe}(\text{H}_2\text{O})_5(\text{NO})]\text{SO}_4$  (B) diamagnetic  $[\text{Fe}(\text{H}_2\text{O})_5(\text{N}_3)]\text{SO}_4$   
 (C) paramagnetic  $[\text{Fe}(\text{H}_2\text{O})_5(\text{NO}_3)](\text{SO}_4)_2$  (D) diamagnetic  $[\text{Fe}(\text{H}_2\text{O})_4(\text{SO}_4)]\text{NO}_3$   
**Solution : (A)**
63. Amongst Be, B, Mg and Al the second ionization potential is maximum for  
 (A) B (B) Be (C) Mg (D) Al  
**Solution : (A)**
64. Sulphuryl chloride ( $\text{SO}_2\text{Cl}_2$ ) reacts with white phosphorus ( $\text{P}_4$ ) to give  
 (A)  $\text{PCl}_5, \text{SO}_2$  (B)  $\text{OPCl}_3, \text{SOCl}_2$  (C)  $\text{PCl}_5, \text{SO}_2, \text{S}_2\text{Cl}_2$  (D)  $\text{OPCl}_3, \text{SO}_2, \text{S}_2\text{Cl}_2$   
**Solution : (A)**
65. The number of lone pair of electrons on the central atoms of  $\text{H}_2\text{O}$ ,  $\text{SnCl}_2$ ,  $\text{PCl}_3$  and  $\text{XeF}_2$  respectively, are  
 (A) 2, 1, 1, 3 (B) 2, 2, 1, 3 (C) 3, 1, 1, 2 (D) 2, 1, 2, 3  
**Solution : (A)**
66. Consider the following salts :  $\text{NaCl}$ ,  $\text{HgCl}_2$ ,  $\text{Hg}_2\text{Cl}_2$ ,  $\text{CuCl}_2$ ,  $\text{CuCl}$  and  $\text{AgCl}$ . Identify the correct set of insoluble salts in water.  
 (A)  $\text{Hg}_2\text{Cl}_2, \text{CuCl}, \text{AgCl}$  (B)  $\text{HgCl}_2, \text{CuCl}, \text{AgCl}$  (C)  $\text{Hg}_2\text{Cl}_2, \text{CuCl}_2, \text{AgCl}$  (D)  $\text{Hg}_2\text{Cl}_2, \text{CuCl}, \text{NaCl}$   
**Solution : (A)**
67. In the following compound, the number of 'sp' hybridized carbon is  

$$\text{CH}_2=\text{C}=\underset{\text{CN}}{\text{CH}}-\text{C}\equiv\text{CH}$$
  
 (A) 2 (B) 3 (C) 4 (D) 5  
**Solution : (C)**
68. Match the flame colours of the alkaline earth metal salts in the Bunsen burner  
 (a) Calcium (p) brick red  
 (b) Strontium (q) apple green  
 (c) Barium (r) crimson  
 (A) a-p, b-r, c-q (B) a-r, b-p, c-q (C) a-q, b-r, c-p (D) a-p, b-q, c-r  
**Solution : (A)**
69. Extraction of gold (Au) involves the formation of complex ions 'X' and 'Y'  

$$\text{Gold ore} \xrightarrow[\text{CN}^-, \text{H}_2\text{O}, \text{O}_2]{\text{Roasting}} \text{HO}^- + \text{'X'} \xrightarrow{\text{Zn}} \text{'Y'} + \text{Au}$$
  
 'X' and 'Y' are respectively  
 (A)  $\text{Au}(\text{CN})_2^-$  and  $\text{Zn}(\text{CN})_4^{2-}$  (B)  $\text{Au}(\text{CN})_4^{3-}$  and  $\text{Zn}(\text{CN})_4^{2-}$   
 (C)  $\text{Au}(\text{CN})_3^-$  and  $\text{Zn}(\text{CN})_6^{4-}$  (D)  $\text{Au}(\text{CN})_4^-$  and  $\text{Zn}(\text{CN})_3^-$   
**Solution : (B)**
70. The atomic number of cerium (Ce) is 58. The correct electronic configuration of  $\text{Ce}^{3+}$  ion is  
 (A)  $[\text{Xe}]4f^1$  (B)  $[\text{Kr}]4f^1$  (C)  $[\text{Xe}]4f^{13}$  (D)  $[\text{Kr}]4d^1$   
**Solution : (A)**




**Category – II (Q71 to Q75)**

Each question has one correct option and carries 2 marks, for each wrong answer 1/2 mark will be deducted.

71. Roasted copper pyrite on smelting with sand produces  
 (A)  $\text{FeSiO}_3$  as fusible slag and  $\text{Cu}_2\text{S}$  as mattee' (B)  $\text{CaSiO}_3$  as infusible slag and  $\text{Cu}_2\text{O}$  as mattee'  
 (C)  $\text{Ca}_3(\text{PO}_4)_2$  as fusible slag and  $\text{Cu}_2\text{S}$  as mattee' (D)  $\text{Fe}_3(\text{PO}_4)_2$  as infusible slag and  $\text{Cu}_2\text{S}$  as mattee'

**Solution : (A)**

72. The total number of aromatic species generated in the following reactions is



- (A) zero (B) 2 (C) 3 (D) 4

**Solution : (C)**

73. For the reaction  $\text{X}_2\text{Y}_4(\text{l}) \rightarrow 2\text{XY}_2(\text{g})$  at 300 K the values of  $\Delta\text{U}$  and  $\Delta\text{S}$  are 2 kCal and  $20 \text{ Cal K}^{-1}$  respectively. The value of  $\Delta\text{G}$  for the reaction is

- (A)  $-3400 \text{ Cal}$  (B)  $3400 \text{ Cal}$  (C)  $-2800 \text{ Cal}$  (D)  $2000 \text{ Cal}$

**Solution : (C)**

74. At temperature of 298 K the emf of the following electrochemical cell



will be (given  $E_{\text{cell}}^0 = -1.562 \text{ V}$ )

- (A)  $-1.532 \text{ V}$  (B)  $-1.503 \text{ V}$  (C)  $1.532 \text{ V}$  (D)  $-3.06 \text{ V}$

**Solution : (A)**

75. Addition of sodium thiosulphate solution to a solution of silver nitrate gives 'X' as white precipitate, insoluble in water but soluble in excess thiosulphate solution to give 'Y'. On boiling in water, 'Y' gives 'Z'. 'X', 'Y' and 'Z' respectively, are


- (A)  $\text{Ag}_2\text{S}_2\text{O}_3$ ,  $\text{Na}_3[\text{Ag}(\text{S}_2\text{O}_3)_2]$ ,  $\text{Ag}_2\text{S}$  (B)  $\text{Ag}_2\text{SO}_4$ ,  $\text{Na}[\text{Ag}(\text{S}_2\text{O}_3)_2]$ ,  $\text{Ag}_2\text{S}_2$   
 (C)  $\text{Ag}_2\text{S}_2\text{O}_3$ ,  $\text{Na}_5[\text{Ag}(\text{S}_2\text{O}_3)_3]$ ,  $\text{AgS}$  (D)  $\text{Ag}_2\text{SO}_3$ ,  $\text{Na}_3[\text{Ag}(\text{S}_2\text{O}_3)_2]$ ,  $\text{Ag}_2\text{O}$

**Solution : (A)**

**Category – III (Q76 to Q80)**

Each question has one or more correct option(s) choosing which will fetch maximum 2 marks on pro rata basis. However, choice of any wrong option(s) will fetch zero mark for the question.

76.







Identify the correct method for the synthesis of the compound shown above from the following alternatives

- (A)  $\xrightarrow[\text{AlCl}_3]{\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{Cl}}$   $\xrightarrow[\text{H}_2\text{SO}_4]{\text{HNO}_3}$
- (B)  $\xrightarrow[\text{AlCl}_3]{\text{CH}_3\text{CH}_2\text{CH}_2\text{COCl}}$   $\xrightarrow[\text{HCl/heat}]{\text{Zn/Hg}}$   $\xrightarrow[\text{H}_2\text{SO}_4]{\text{HNO}_3}$
- (C)  $\xrightarrow[\text{AlCl}_3]{\text{CH}_3\text{CH}_2\text{CH}_2\text{COCl}}$   $\xrightarrow[\text{H}_2\text{SO}_4]{\text{HNO}_3}$   $\xrightarrow[\text{HCl/heat}]{\text{Zn/Hg}}$
- (D)  $\xrightarrow[\text{AlCl}_3]{\text{CH}_3\text{CH}_2\text{CH}_2\text{COCl}}$   $\xrightarrow[\text{OH}^-]{\text{KMnO}_4}$   $\xrightarrow[\text{H}_2\text{SO}_4]{\text{HNO}_3}$

**Solution : (B)**

77. Within the list shown below, the correct pair of structures of alanine in pH ranges 2-4 and 9-11 is



(A) I, II

(B) I, III

(C) II, III

(D) III, IV

**Solution : (A)**

78. The increase in rate constant of a chemical reaction with increasing temperature is(are) due to the fact(s) that

(A) the number of collisions among the reactant molecules increases with increasing temperature.



(B) the activation energy of the reaction decreases with increasing temperature.

(C) the concentration of the reactant molecules increases with increasing temperature.

(D) the number of reactant molecules acquiring the activation energy increases with increasing temperature.

**Solution : (A, C, D)**

79. Optical isomerism is exhibited by (ox = oxalate anion ; en = ethylenediamine)



**Solution : (A, B, D)**



80. Ionization potential values of noble gases decrease down the group with increase in atomic size. Xenon forms binary fluorides by the direct reaction of elements. Identify the correct statement(s) from below.
- (A) Only the heavier noble gases form such compounds.
  - (B) it happens because the noble gases have higher ionization energies.
  - (C) It happens because the compounds are formed with electronegative ligands.
  - (D) Octet of electrons provide the stable arrangements.

**Solution :** (A, C)

**END OF QUESTION PAPER**

