Total Questions: 200

Max. Time: 3 1/2 hrs.

Instructions

- Each question has four options. Mark the most appropriate choice as correct answer.
- Follow the instructions for answering, given on the OMR sheet and BACK page of Admit Card.
- Enter your Roll Number, Serial No. of OMR answer sheet, full signature and name in the spaces provided on the question paper, duplicate face sheet, answer sheet and attendance list.
- Use only blue or black ball point pen for writing/marking
- For each correct answer, one mark will be awarded. Each incorrect answer will be awarded

one third $\left(-\frac{1}{3}\right)$ negative mark. **Zero Mark** will be

given for question not answered. Any answer having more than one entry, will be treated as wrong answer and awarded negative mark.

- Do not mark or write anything on the question booklet
- Any candidate found removing page(s) and/or copying down questions or using unfair means will be disqualified. Pager, cellular phone, calculator or any such electronic devices are strictly prohibited in the examination hall.
- Any discrepancy or ambiguity in any question may be reported to Sub - Dean (Exams) in writing within 72 hours. No notice will be taken of representations received after 72 hours
- No candidate will be allowed to leave the Hall until
 - Three and half hours have elapsed after the start of the Examination
 - (2) The answer sheet is countersigned by both the Invigilators
 - (3) The Invigilator in your half has taken your signature in the attendance list.
- § In the following questions (1 60), a statement of assertion (1) is followed by a statement of reason (R)

If both Assertion and Reason are true and the reason is the correct explanation of the assertion, then mark 1.

If both Assertion and Reason are true but the reason is not the correct explanation of the assertion, then mark 2.

If Assertion is true statement but Reason is false, then mark 3.

If both Assertion and Reason are false statements, then mark 4.

- A : A person who has received a cut and is bleeding needs to be given anti-tetanus treatment.
 - R : Anti-tetanus injection provides immunity by producing antibodies for tetanus
- A : Cancer cells are virtually immortal until the body in which they reside dies.
 - R : Cancer is caused by damage to genes regulating the cell division cycle
- A : A network of food chains existing together in an ecosystem is known as a food web.
 - R : An animal like kite cannot be a part of a food web.
- A: Inflammation of a skeletal joint may immobilize the movements of the joint

- R: Uric acid crystals in the joint cavity and ossification of articular cartilage lead to this.
- A : The earliest organisms that appeared on the Earth were non-green and presumably anaerobes.
 - R : The first autotrophic organisms were the chemoautotrophs that never released oxygen.
- A : Escherichia coli, Shigella sp. and Salmonella sp. are all responsible for diarrhoeal diseases.
 - R : Dehydration is common to all types of diarrhoeal diseases and adequate supply of fluids and electro lytes should be ensured.
- A : Deforestation is one main factor contributing to global warming
 - R : Besides CO₂, two other gases methane and CFCs are also included under green house gases
- CFCs are also included under green house gases.

 8. A : LSD and marijuana are clinically used as anal-
 - R : Both these drugs suppress brain function
- A: An organism with lethal mutation may not even develop beyond the zygote stage
- R : All types of gene mutations are lethal
- 10. A : Our body secretes adrenaline in intense cold
 - R : Adrenaline raises metabolic rate
- A : In collateral vascular-bundles, phloem is situated towards inner side
- R: In monocot stem, cambium is present.
- A: In angiosperms the conduction of water is more efficient because their xylem has vessels

- R : Conduction of water by vessel elements is an active process with energy supplied by xylem parenchyma rich in mitochondria
- A : Polytene chromosomes have a high amount of DNA
 - R : Polytene chromosomes are formed by repeated replication of chromosomal DNA without separation of chromatids
- A: UV radiation causes photo dissociation of ozone into O₂ and O, thus causing damage to the stratospheric ozone layer.
 - R : Ozone hole is resulting in global warming and climate change
- A : The concentration of methane in the atmosphere has more than doubled in the last 250 years.
 - R : Wetlands and rice fields are the major sources of methane
- A: In tropical rain forests, O-Horizon and A-Horizon of soil profile are shallow and nutrient-poor.
 - R : Excessive growth of micro-organisms in the soil depletes its organic content.
- A: Gram-negative bacteria do not retain the stain when washed with alcohol
 - R : The outer face of the outer membrane of Gram-negative bacteria contains lipopolysaccharides, a part of which is integrated into the membrane lipids.
- 18. A: Under conditions of high intensity and limited CO₂ supply, photo respiration has a useful role in protecting the plants from photo-oxidative damage. R: If enough CO₂ is not available to utilize light energy for carboxylation to proceed, the excess energy may not cause damage to plants.
- A: Photo synthetically C₄ plants are less efficient than C₃ plants
 - R : The operation of C₄ pathway requires the involvement of only bundle-sheath cells.
- 20. A : Eukaryotic cells have the ability to adopt a variety of shapes and carry out directed movements. R : There are three principal types of protein filaments- microfilaments, microtubules and intermediate filaments, which constitute the cytoskeleton.
- A : In the iodometric titration starch is used as an indicator
 - R : Starch is polysaccharide
- A : Molecular nitrogen is less reactive than molecular oxygen
 - R : The bond length of N₂ is shorter than that of oxygen
- A : [Co(NO₂)₃ (NH₃)₃] does not show optical isomerism
 - R: It has a plane of symmetry.
- 24. A : E° for Mn³+/Mn²+ is more positive than

 Cr³+/Cr²-

- R: The third ionization energy of Mn is larger than that of Cr
- A: K₂Cr₂O₇ is used as a primary standard in volumetric analysis
 - R : It has a good solubility in water
- 26. A : Silicones are hydrophobic in nature
 - R: Si O Si linkages are moisture sensitive
- 27. A : According to transition state theory, for the formation of an activated complex, one of the vibrational degree of freedom is converted into a translational degree of freedom
 - R: Energy of the activated complex is higher than the energy or reactant molecules
- 28. A : Water in liquid state is more stable than ice at room temperature
 R : Water in liquid from has higher entropy than
- ice. 29. A: Sb₂ S₃ is not soluble in yellow ammonium sul-
- phide
 - ${f R}$: The common ion effect due to $S^{2^{n}}$ ions reduces the solubility of $Sb_2\,S_3$
- A : Graphite is an example of tetragonal crystal system
 - R : For a tetragonal system
 - $a = b \neq c$, $\alpha = \beta = 90^{\circ}$, $\gamma = 120^{\circ}$
- A : For the Daniel cell Zn|Zn²+||Cu²+|Cu with E_{cell} = 1.1 V, the application of opposite potential greater than 1.1 V results into flow of electron from cathode to anode
 - R : Zn is deposited at anode and Cu is dissolved at cathode
- 32. A : Fe3+ can be used for coagulation of As₂S₃ sol.
 - R : Fe3+ reacts with As2S3 to give Fe2S3
- A: If red blood cells were removed from the body and placed in pure water, pressure inside the cells increases
 - R : The concentration of salt content in the cells in-
- 34. A : Change in colour of acidic solution of potassium dichromate by breath is used to test drunk drivers.
 - R : Change in colour is due to the complication of alcohol with potassium dichromate
- A : Anilinium chloride is more acidic than ammonium chloride
 - R : Antlinium ion is resonance-stabilized
- A : Diastereoisomers have different physical properties
 - R: They are non-superimposable mirror images.
- A: The presence of nitro group facilitates nucleophilic substitution reactions in aryl halides
 - R : The intermediate carbanion is stabilized due to the presence of nitro group
- 38 A : 1, 3-Butadiene is the monomer for natural rubber

R: Natural rubber is formed through anionic addi- 51. A: A man in a closed cabin falling freely does not tion polymerization

39. A : Addition of HBr on 2-butene gives two isomeric products

R .: Addition of HBr on 2-butene follows Markovnikov rule

40. A : The water pouch of instant cold pack for treating athletic injuries breaks when squeezed and NH4NO3 dissolves lowering the temperature

R : Addition of non-volatile solute into solvent results into depression of freezing point of the solvent.

41. A : Electromagnetic waves with frequencies smaller than the critical frequency of ionosphere cannot be used for communication using sky wave propagation

R : The refractive index of the ionosphere becomes very high for frequencies higher than the critical frequency

- 42. A : The binding energy per nucleon, for nuclei with atomic mass number A > 100, decreases with
 - R : The nuclear forces are weak for heavier nuclei
- 43. A : In common base configuration, the current gain of the transistor is less than unity

R: The collector terminal is reverse biased for amplification

- 44. A : In an isolated system the entropy increases R : The processes in an isolated system are adiabatic
- 45. A : Magnetic Resonance Imaging (MRI) is a useful diagnostic tool for producing images of various parts of human body.

R : Protons of various tissues of the human body play a role in MRI

46. A : A judo fighter in order to throw his opponent on to the mat tries to initially bend his opponent and them rotate him around his hip

R : As the mass of the opponent is brought closer. to the fighter's hip, the force required to throw the opponent is reduced

47. A : The root mean square and most probable speeds of the molecules in a gas are the same

R : The Maxwell distribution for the speed of molecules in a gas is symmetrical

48. A : Use of ball bearings between two moving parts of a machine is a common practice

R : Ball bearings reduce vibrations and provide good stability.

49. A : Standard optical diffraction gratings can not be used for discriminating between different X-ray wavelengths

R : The grating spacing is not of the order of X-ray wavelengths

 A: Diamagnetic materials can exhibit magnetism R : Diamagnetic materials have permanent magnetic dipole moment

experience gravity

R : Inertial and gravitational mass have equiva-

52. A : The photoelectrons produced by a monochromatic light beam incident on a metal surface, have a spread in their kinetic energies

R : The work function of the metal varies as a function of depth from the surface

53. A : The Carnot cycle is useful in understanding the performance of heat engines R : The Carnot cycle provides a way of determining the maximum possible efficiency achievable

54. A : A p-n junction with reverse bias can be used as a photo-diode to measure light intensity

with reservoirs of given temperatures

R : In a reverse bias condition the current is small but it is more sensitive to changes in incident light

55. A : Perspiration from human body helps in cooling the body

R : A thin layer of water on the skin enhances its emissivity

56. A: When a glass of hot milk is placed in a room and allowed to cool, its entropy decreases R : Allowing hot object to cool does not violate the

57. A : Cobalt-60 is useful in cancer therapy

second law of thermodynamics

R : Cobalt-60 is a source of y-radiations capable of killing cancerous cells.

58. A : A thin stainless steel needle can lay floating on a still water surface.

R : Any object floats when the buoyancy force balances the weight of the object

59. A : An emf E is induced in a closed loop where magnetic flux is varied. The induced E is not a conservative field

R : The line integral E . at around the closed loop is non-zero

60.A : In optical fibre, the diameter of the core is kept small

R: This smaller diameter of the core ensures that the fibre should have incident angle more than the critical angle required for total internal reflection

61. The pair in which both species have same magnetic moment (spin only value) is :

(1) [Cr(H2O)6]2+, [CoCl4]2-

(2) [Cr(H2O)6]2+, [Fe(H2O)6]2+

(3) [Mn(H2O)6]2+, [Cr(H2O)6]2+

(4) [CoCl₄]2-, [Fe(H₂O)₆]2+

62. The pair in which both species have iron is :

Nitrogenase, cytochromes

(2) Carboxypeptidase, Haemoglobin

- (3) Haemocyanin, nitrogenase
- (4) Haemoglobin, cytochromes
- 63. Borax is used as cleaning agent because on dissolving in water it gives :
 - (1) Alkaline solution
 - (2) Acidic solution
- (3) Bleaching solution
- (4) Colloidal solution
- 64. The incorrect statement among the following is :
 - (1) C₆₀ is an allotropic form of carbon
 - (2) O₃ is an allotropic form of oxygen
 - (3) Se is only allotropic form of sulphur
 - (4) Red phosphorus is more stable in air than white phosphorus
- 65. The pair whose both species are used in anti acid medicinal preparations is :
 - (1) NaHCO3 and Mg(OH)2
 - (2) Na₂CO₃ and Ca(HCO₃)₂
 - (3) Ca(HCO₂)₂ and Mg(OH)₂
 - (4) Ca(OH)₂ and NaHCO₃
- 66. The colour imparted by Co(II) compounds to glass is :
 - (1) Green
- (2) Deep -Blue
- (3) Yellow
- (4) Red
- 67. The ligands in anti cancer drug cisplatin are:
 - (1) NH2, CI
- (2) NH₃, H₂O
- (3) Cl. H₀O
- (4) NO. CI
- 68. The number of possible isomers of an octahedral complex [Co(C2O4)2 (NH3)2]" is :
 - (2)2(3) 3
- 69. Given below, catalyst and corresponding process/reaction are matched. The mismatch is :
 - (1) [RhCl(PPh₃)₂] : Hydrogenation
 - (2) TiCl₄ + Al(C₂H₅)₃: Polymerization
 - (3) V₂O₅: Haber-Bosch process
 - (4) Nickel: Hydrogenation
- 70. Among the following, the species having square planar geometry for central atom are
 - (i) XeF4 (ii) SF4 (iii) [NICL]2- (iv) [PdCL]2-
 - (1) (i) and (iv) (3) (ii) and (iii)
- (2) (i) and (ii)
- (4) (iii) and (iv)
- 71. Tincture of iodine is:
 - (1) Aqueous solution of la
 - (2) Solution of I2 in aqueous KI
 - (3) Alcoholic solution of I,
 - (4) Agueous solution of KI

- 72. In [Ag(CN)₂], the number of π bonds is :
 - (2) 3
- (3)4
- 73. The compound molecular in nature in gas phase but ionic in solid state is :
 - (2) CCL
- (3) PC/2
- 74. Which two of the following salts are used for preparing iodized salt?
 - (i) KIO₃ (ii) KI (iii) I₂ (iv) HI:
 - (1) (i) and (ii)
- (2) (i) and (iii)
- (3) (ii) and (iv)
- (4) (iii) and (iv)
- 75. The compound used in enrichment of uranium for nuclear power plant is :
 - (1) U₂O₆
- (2) UF4
- (3) UO2(NO2)2
- (4) UCL
- 76. The de Broglie wavelength associated with a ball of mass 1 kg having kinetic energy 0.5
 - (1) 6.626×10^{-34} m (2) 13.20×10^{-34} m
 - (3) 10.38 × 10⁻²¹ m (4) 6.626 × 10⁻³⁴ Å
- 77. Dominance of strong repulsive forces among the molecules of the gas (Z = compressibility factor):
- (1) Depends on Z and indicated by Z = 1
 - (2) Depends on Z and indicated by Z > 1
 - (3) Depends on Z and indicated by Z < 1</p>
 - (4) Is independent of Z
- 78. 40 ml of 0.1 M ammonia solution is mixed with 20 ml of 0.1 M HCl. What is the PH of the mixture ? (pk. of ammonia solution is 4.74)
 - (1) 4.74
- (2)2.26
- (3)9.26
- (4)5.00
- 79. For a spontaneous process the correct statement is :
 - (1) Entropy of the system always increases
 - (2) Free energy of the system always increases
 - (3) Total entropy change is always negative
 - (4) Total entropy change is always positive
- 80. The Ca2+ and F are located in CaF, crystal, respectively at face centred cubic lattice points and in
 - (1) Tetrahedral voids (2) Half of tetrahedral voids
- (3) Octahedral voids (4) Half of octahedral voids 81. The charge required for the reduction of 1
 - mol of MnO, to MnO, is:
 - (1) 1 F (2) 3 F
- (3) 5 F
- 82. For the reaction

2 N,O, --- 4 NO, + O, rate of reaction is :

(1) $\frac{1}{2} \frac{d}{dt} [N_2O_5]$ (2) $2 \frac{d}{dt} [N_2O_5]$

(3) $\frac{1}{4} \frac{d}{dt} [NO_2]$ (4) $4 \frac{d}{dt} [NO_2]$

83. For a phase change

(1) $\Delta G = 0$

(2) $\Delta S = 0$

(3) $\Delta H = 0$

(4) AU = 0

- 84. A 5% solution (by mass) of cane sugar in water has freezing point of 271 K and freezing point of pure water is 273.15 K. The freezing point of a 5 % solution (by mass) of glucose in water is
 - (1) 271 K

(2) 273.15 K

(3) 269.07 K

(4) 277.23 K

- 85. The energy gaps (E,) between valence band and conduction band for diamond, silicon and germanium are in the order
 - (1) E_n (diamond) > E_n (silicon) > E_n (germa-
 - (2) E (diamond) < E (silicon) < E (germanium)
 - (3) En (diamond) = En (silicon) = En (germanium)
 - (4) E (diamond) > E (germanium) > E (silicon)
- 86. The enthalpy change (AH) for the reaction, $N_2(g) + 3 H_2(g) \rightarrow 2 NH_3(g)$ is - 92.38 kJ at 298 K. The internal energy change ∆U at 298 K is

(1) - 92.38 W

(2) - 87.42 kJ

(3) - 97.34 kJ

(4) - 89.9 kJ

- 87. The products formed when an aqueous solution of NaBr is electrolyzed in a cell having inert electrodes are :
 - (1) Na and Bra

(2) Na and O2


(3) H2, Br2 and NaOH

(4) H2 and O2

88. Among the following L-series is

- 89. Among the following which one can have a meso form ?
 - CH₃CH(OH)CH(CI)C₂H₅
 - (2) CH₃CH(OH)CH(OH)CH₃
 - (3) C.H.CH(OH)CH(OH)CH.
 - (4) HOCH₂CH(CI)CH₃
- 90. Which of the following sequence of reactions (reagents) can be used for the conversion of CaHaCHaCHa in to CaHaCH = CHa?
 - (1) SOCI.; HO
 - (2) SO₂Cl₂: alc. KOH
 - (3) Cl2/hv; H2O
 - (4) SOCIo: alc. KOH
- 91. Isopropylbenzene on air oxidation in the presence of dilute acid gives :
 - · (1) C,H,COOH
 - (2) C_cH_cCOCH₂
 - (3) C₄H₅CHO
 - (4) C.H.OH
- 92. Nitrobenzene on treatment with zinc dust and aqueous ammonium chloride gives :
 - (1) C. H. N = N C. H.
 - (2) C₆H₆NH₂
 - (3) C. H. NO
 - (4) Cs Hs NHOH
- 93. The major product obtained on the monobromination (with Br2 /FeBr3) of the following compound A is

94. The following sequence of reactions on A gives



- 95. Thymine is :
 - (1) 5 Methyluracil
- (2) 4 Methyluracil
- (3) 3 Methyluracil
- (4) 1 Methyluracil 96. Lysine is least soluble in water in the pH
- range :
 - (1) 3 to 4
- (2),5 to 6
- (3) 6 to 7
- (4) 8 to 9
- 97. Methyl $-\alpha D$ glucoside and methyl $-\beta$ -D - Glucoside are :
 - (1) Epimers
- (2) Anomers
- (3) Enantiomers
- (4) Conformational diastereomers
- 98. Which of the following compounds has the highest boiling point?
 - (1) CH, CH, CH, CI (2) CH, CH, CH, CH, CH
 - (3) CH₃ CH (CH₃) CH₂ CI
 - (4) (CH₂)₂ C C/
- 99. The correct increasing order of the reactivity of halides for SNI reaction is :
 - (1) CH₃ CH₂ X < (CH₃)₂ CH X < CH2 = CH - CH2 - X < PhCH2 - X
 - (2) (CH₃)₂CH X < CH₃ CH₂ X <

$$CH_2 = CH - CH_2 X < PhCH_2 - X$$

(3) PhCH₂ - X < (CH₂)₂CH - X < CH₃ - CH₃ - X < CH₂ = CH - CH₃ - X

100. The major product formed in the following reaction :

- (1) CH, CH = CH CH, OH
- (2) CH2 = CH CH2 CH2OH

- 101. In refraction, light waves are bent on passing from one medium to the second medium, because, in the second medium.
 - (1) The frequency is different
 - (2) The coefficient of elasticity is different
 - (3) The speed is different
 - (4) The amplitude is smaller
- 102. Two spheres of same size, one of mass 2 Kg and another of mass 4 Kg, are dropped simultaneously from the top of Qutab Minar (height = 72 m). When they are 1 m above the ground, the two spheres have the same:
 - (1) Momentum
- (2) Kinetic energy
- (3) Potential energy (4) Acceleration
- 103. The moment of inertia of a rod about an axis through its centre and perpendicular to it is $\frac{1}{12}$ M L² (where M is the mass and L, the


length of the rod). The rod is bent in the middle so that the two halts make an angle of 60°. The moment of inertia of the bent rod about the same axis would be :

- (1) 1 ML2 ·
- (2) 12 M L2
- (3) 1 M L2
- (4) ML²
- 104. A boat at anchor is rocked by waves whose crests are 100 m apart and velocity is 25 m/ sec. The boat bounces up once in every
 - (1) 2500 s
- (2) 75 s
- (3) 4 s
- (4) 0.25 s
- 105. By sucking through a straw, a student can reduce the pressure in his lungs to 750 mm of Hg (density 13.6 gm/ cm3). Using the straw, he can drink water from a glass upto a maximum depth of :
 - (1) 10 cm
- (2) 75 cm
- (3) 13.6 cm
- (4) 1.36 cm
- 106. Two parallel large thin metal sheets have surface charged $(\sigma = 26.4 \times 10^{-12} \text{ C/m}^2)$ of opposite signs. The electric field between these sheets is :
 - (1) 1.5 N/C
- (2) 1.5 × 10⁻¹⁰ N/C
- (3) 3 N/C
- (4) 3 x 10⁻¹⁰ N/C
- 107. The magnetic moment has dimensions of :
 - (1) [L, A]
- (2) (L2 A)

- (4) (L2 T-1 A)
- (3) [L T · 1 A] (4) [L² T · 1 A] 108. A wire mesh consisting of very small squares is viewed at a distance of 8 cm through a magnifying converging lens of focal length 10 cm, kept close to the eye. The magnification produced by the lens is :
 - (1)5

- (2) 8
- (3) 10


- (4)20
- 109. Hard X-rays for the study of fractures in bones should have a minimum wavelength of 10-11 m. The accelerating voltage for electrons in X-ray machine should be
 - (!) < 124.2 kV
 - (2) > 124.2 kV
 - (3) Between 60 kV and 70 kV
 - (4) = 100 kV
- 110. In photoelectric effect, the electrons are ejected from metals if the incident light has a certain minimum :
 - (1) Wavelength
- (2) Frequency
- (3) Amplitude
- (4) Angle of incidence
- 111. A lens is made of flint glass (refractive index = 1.5). When the lens is immersed in a liquid of refractive index 1.25, the focal length:
 - (1) increases by a factor of 1.25
 - (2) increases by a factor of 2.5
 - (3) increases by a factor of 1.2
 - (4) decreases by a factor of 1.2
- 112. The voltage of clouds is 4×10^6 volt with respect to ground. In a lightening strike lasting 100 m sec, charge of 4 coulombs is delivered to the ground. The power of lightening strike is :
 - (1) 160 MW
- (2) 80 MW
- (3) 20 MW
- (4) 500 KW
- 113. Five capacitors, each of capacitance value C are connected as shown in the figure. The ratio of capacitance between P and R, and the capacitance between P and Q is :




- (1) 3:1
- (2) 5:2
- (3)2:3
- (4) 1:1

- 114. A stone thrown into still water creates a circular wave pattern moving radially outwards. If r is the distance measured from the centre of the pattern, the amplitude of the wave varies as :
 - (1) +1/2 (3) r2
- 115. For inelastic collision between two spherical rigid bodies :
 - (1) The total kinetic energy is conserved
 - (2) The total potential energy is conserved
 - (3) The linear momentum is not conserved
 - (4) The linear momentum is conserved
- 116. When a p-n diode is reverse biased, then :
 - (1) No current flows
 - (2) The depletion region is increased
 - (3) The depletion region is reduced
 - (4) The height of the potential barrier is reduced
- 117. The operation of a nuclear reactor is said toe be critical, if the multiplication factor (k) has a value
 - (1) 1


- (2)1.5
- (3)2.1
- (4)2.5
- 118. Which of the following diagrams represent the variation of electric field vector with time for a circularly polarized light?




119. The circuit given below representes which of logic operations :




- (I) AND (2) NOT
- (3) OR
- (4) NOR
- 120. A person used force (F), shown in figure to move a load with constant velocity on given surface




Identify the correct surface profile




- 121. Three objects coloured black, gray and white can withstand hostile conditions upto 2800°C. These objects are thrown into a furnace where each of them attains a temperature of 2000°C. Which object will glow brightest?
 - (1) the white object (2) the black object
 - (3) all glow with equal brightness
 - (4) gray object
- 122. Two balloons are filled, one with pure He gas and the other by air, respectively. If the pressure and temperature of these balloons are same then the number of molecules per unit volume is :
 - (1) More in the He filled balloon
 - (2) Same in both balloons
 - (3) More in air filled balloon
 - (4) In the ratio of 1 : 4
- 123. Flash light equipped with a new set of batteries, products bright white light. As the batteries wear out :
 - The light intensity gets reduced with no change in its colour
 - (2) Light colour changes first to yellow and then red with no change in intensity
 - (3) It stops working suddenly while giving white light
 - (4) Colour changes to red and also intensity gets reduced
- 124. The spatial distribution of the electric field due to two charges (A, B) is shown in figure. Which one of the following statements is correct



- (1) A is + ve and B ve and |A| > |B|
- (2) A is ve and B + ve ; |A| = |B|
- (3) Both are +ve but A> B
- (4) Both are ve but A > B 125. Circular loop of a wire and a long straight wire carry currents Ic and I. . respectively shown in figure. Assuming that these are placed in the same plane. the magnetic fields will be zero at the centre of the loop when the separation H is :



- $(2) \frac{L_R}{L_R} \qquad (3) \frac{\pi L}{L_R} \qquad (4) \frac{L_R}{L_R}$
- 126. If a street light of mass M is suspended from the end of a uniform rod of length L in different possible patterns as shown in figure



- Pattern A is more sturdy
- (2) Pattern B is more sturdy
- (3) Pattern C is more sturdy
- (4) All will have same sturdiness
- 127. 238U.has 92 protons and 238 nucleons. It decays by emitting an Alpha particle and becomes:
 - (2) 234Th (3) 235U (4) 237Np (1) 234U
- 128. The fossil bone has a 14C: 12C ratio. which is $\left(\frac{1}{16}\right)$ of that in a living animal bone.

If the half-life time of 14C is 5730 years, then the age of the fossil bone is :

- (1) 11460 years
- (2) 17190 years

(3) 22920 years

(4) 45840 years

129. Which one of the following is a possible nuclear reaction :

(3)
$$^{239}_{93}Np \longrightarrow ^{239}_{94}Pu + \beta^- + \nu^-$$

$$(4)_{7}^{11}N + {}_{1}^{1}H \longrightarrow {}_{6}^{12}C + \beta^{-} + v$$

130. When a guitar string is sounded with a 440 Hz tuning fork, a beat frequency of 5 Hz is heard. If the experiment is repeated with a tuning fork of 437 Hz, the beat frequency is 8 Hz. The string frequency (Hz) is :


(1)445

(2) 435

(3) 429

(4)448

131. A metallic ring is dropped down, keeping its plane perpendicular to a constant and horizontal magnetic field. The ring enters the region of magnetic field at t=0 and completely emerges out at t=T sec. The current in the ring varies as :




- 132. If Alpha, Beta and Gamma rays carry same momentum, which has the longest wavelength:
 - (1) Alpha rays
- (2) Beta rays
- (3) Gamma rays
- (4) None, all have some wavelength
- 133. An amplifier has a voltage gain A, = 1000, The voltage gain in dB is :
 - (1) 30 dB
- (2) 60 dB
- (3) 3 dB
- (4) 20 dB
- 134. When you make ice cubes, the entropy of water:
 - (1) Does not change
 - (2) Increases

- (3) Decreases
- (4) May either increase or decrease depending on the process used
- 135. A bimetallic strip consists of metals X and Y. It is mounted rigidly at the base as shown. The metal X has a higher coefficient of expansion compared to that for metal Y. When the bimetallic strip is placed in a cold bath :



- (1) It will bend towards the right
- (2) It will bend towards the left
- (3) It will not bend but shrink
- (4) It will neither bend nor shrink
- 136. For a wave propagating in a medium, identify the property that is independent of the others :
 - (1) Velocity
- (2) Wavelength
 - (3) Frequency
 - (4) All these depend on each other
- 137. A leaf which contains only green pigments, is illuminated by a laser light of wavelength 0.6328 µm. It wound appear to be :
 - (1) Brown
- (2) Black
- (3) Red
- (4) Green
- 138. A light emitting diode (LED) has a voltage drop of 2 volt across it and passes a current of 10 mA. When it operates with a 6 volt battery through a limiting resistor R. The value of R is:
 - (1) $40 \, k\Omega$
- (2) 4 kΩ
- (3) 200Ω
- (4) 400Ω
- 139. The minimum potential difference between the base and emitter required to switch a silicon transistor 'ON' is approximately :
 - (1) 1 V
- (2) 3 V
- (3) 5 V
- (4) 4.2 V
- 140. Given below is the circuit diagram of an AM demodulator



For good demodulation of AM signal of car- 149, X-rays were discovered by : rier frequency f, the value of RC should be :

- (1) RC = 1/f
- (2) RC < 1/f
- (3) RC ≥ 1/f
- (4) RC > > 1/f

141. The term 'CPR' often used in first aid stands for :

- (1) Clinic for Pulmonary Rehabilitation
- (2) Chemical Prevention of Rhinitis
- (3) Chemo-prophylaxis response
- (4) Cardio-pulmonary resuscitation

142. Alzheimer's Disease affects:

- (1) Childhood
- (2) Adolescent
- (3) Young people (4) Elderly people
- 143. Central Council for Research in Yoga and Naturopathy (CCRYN) is an organization that funds research in the given field in our country. It is located in :
 - (1) New Delhi
- (2) Rishikesh
- (3) Bangalore
- (4) Huderabad
- 144. The vaccination for which one of the following diseases is not covered in the immunization schedule so far ?
 - (1) Tuberculosis
- (2) Diphtheria
- (3) Measles
- (4) Pneumonia

145. Which one of the following is true and is not a misconception?

- (1) Cutting the top of a bitter cucumber and rubbing it with the corresponding surface removes
- (2) Pearl is produced from a drop of rain water falling into the cyster in Swati Nakshatra
- (3) While dreaming during sleep at night the eyes. move rapidly
- (4) Peahen gets conceived from the tears the drop from the peacock's eye and she drinks them.

146. A drug called reserpine was discovered by:

- (1) Jal Vakil
- (2) Paul Ehrlish
- (3) Hansen
- (4) Alexander Wood

147. Biometry refers to :

- (1) Identification of humans by scanning face and fingerprints
- (2) Measurement of mechanical displacement in humans
- (3) A method of lie detection
- (4) Body length relationships across the evolutionary scale
- 148. Which one of the following is one of the two days when the sun rises exactly in the enst?
 - (!) 14th January
- (2) 21st March
- (3) 21st June
- (4) 23rd December

- (1) Wilhelm K. Roentgen
- (2) H. Kissinger
- (3) Sir C. V. Raman (4) Meghnad Saha
- 150. Which one of the following literary titles is correctly matched with its author?
 - (1) Ramayana
- Tulsidas
- (2) Mahabharat (3) Kumarsambhav
- Vedvuas - Ravidas
- (4) Shakuntala
- Bhushan
- 151. A very much publicized treatment method "DOTS" is being adopted for the cure of : (3) Tuberculosis
 - (1) Dimentia (2) Tetanus
 - (4) Sexually transmitted disease
- 152. Which of the following Indian cricket player after India-Pakistan ODI (one-day International) at Abudhabi became no. 1 ODI batsman in the ICC (International Cricket Club) ranking:
 - (1) Rahul Dravid
- (2) Yuvrai
- (3) Sachin Tendulkar (4) M. S. Dhoni
- 153. The Jungle in Rudyard Kipplling's Jungle book, describes which part of Indian forest?
- Central Indian forest near Satpura range
 - (2) Uttranchal thick forest
 - (3) Himalayan forest in Himachal
 - (4) Niligiri Jungles

154. Sardar Sarovar Dam is built on the river (1) Jhelam (2) Narmada

- (3) Tapti
- (4) Was
- 155. One ream of paper equals to :
 - (1) 100 110 sheets (2) 256 sheets
 - (3) 480 500 sheets (4) 1000 sheets
- 156. Which of the following honour is given by UNESCO?
 - (1) The Kalinga prize (2) Magasay Award
 - (3) Pulitzer Prize
 - (4) Order of the Golden Ark Award

157. 'Body line' in cricket refers to

- Bowling that hits the body
- (2) The line of body close to Wicket line
- (3) The white line on ground within which the player stands
- (4) The line of moving ball

158. 'Hindu view of life' is written by :

- (1) S. Radhakrishnan (2) R. K. Narayan
- (3) V. D. Savarkar (4) John Ruskin

159. Lagoon refers to:

- (1) A full moon
- ote: The Information provided here is only for reference .It may vary the Original

- (2) The sea breaking into the land and then separated by sand dunes
- (3) A spot in a desert made fertile by presence of water
- (4) Horse shoe shaped coral reef.

160. Ecology deals with :

- (1) The earth and planets
- (2) The relationship between organism and their environments
- (3) The life under the sea
- (4) Economical growth of poor people

161. Given below is a table comparing the effects of sympathetic and parasympathetic nervous system for four features (1-4). Which one feature is correctly described?

	Feature	Sympathetic NervousSystem	Parasympathetic NervousSystem
1.	Salivary glands	Stimulates secretion	Inhibits secretion
2.	Pupil of the eye	Dilate	Constricts
3.	Heart rate	Decreases	Increases
4.	Intestinal peristalsis	Stimulates	Inhibits

162. Which one of the following animals is correctly matched with its one characteristic and the terror.

	Animat	Characteristic	Taxon
1.	Millipede	Ventral nerve cord	Arachnida
2.	Duckbill platypus	Oviparous	Mammalia
3.	Silver fish	Pectoral and Pelvic fins	Chordata
4.	Sea anemone	Triploblastic	Cnidaria

- 163. All mammals without any exception are characterized by :
 - (1) Viviparity and biconcave red blood cells
 - (2) Extra-abdominal testes and a four chambered heart
 - (3) Heterodont teeth and 12 pairs of cranial nerves
 - (4) A muscular diaphragm and milk producing glands
- 164. Which one of the following pairs of the kind of cells and their secretion is correctly matched?
 - Oxyritic cells a secretion with pH between 2.0 and 3.0
 - (2) Alpha cells of Islets of Langerhans secretion that decreases blood sugar level
 - (3) Kupfer cells a digestive enzyme that hydrolyses nucleic acids
 - Sebaceous glands a secretion that evaporates for cooling


- 165. Examine the diagram of the two cell types A and B given below and select the correct option:
 - Cell A is the rod cell found evenly all over retina
 - (2) Cell A is the cone cell more concentrated in the fovea centralis
 - (3) Cell B is concerned with colour vision in bright light
 - (4) Cell A is sensitive to low light intensities
- 166. Which one of the following pairs of features is a good example of polygenic inheritance?
 - (1) Human height and skin colour
 - (2) ABO blood group in humans and flower colour of Mirabilis jalapa
 - (3) Hair pigment of mouse and tongue rolling in humans
- (4) Human eye colour and sickle cell anaemia
- 167. Mating of an organism to a double recessive in order to determine whether it is homozygous or heterozygous for a character under consideration is called:
 - (1) Reciprocal cross (2) Test cross
 - (3) Dihybrid cross (4) Back cross
- 168. In which one of the following sets of three items each belong to the category mentioned against them:
 - (1) Lysine, glysine, thiamine Amino acids
 - (2) Myosin, oxytocin and gastrin Hormones
 - (3) Rennin, helicase and hyaluronidase Enzymes
 - (4) Optic nerve, oculomotor, vagus Sensory nerves
- 169. A cricket player is fast chasing a ball in the field. Which one of the following groups. of bones are directly contributing in this movement?
 - (1) Femur, malleus, tibia, metatarsals
 - (2) Pelvis, ulna, patella, tarsals
 - (3) Sternum, femur, tibia, fibula
 - (4) Tarsals, femur, metatarsals, tibia
- 170. Given: a highly simplified representation of the human sex chromosomes from a karyotype:

The genes a and b could be of:

- colour blindness and body height
- (2) attached ear lobe and Rhesus blood group



- (3) haemophilia and red-green colour blindness
- (4) Phenylketonuria and haemophilia
- 171. A lizard-like member of Reptilia is sitting on a tree with its tall coiled around a twig. This animal could be:
 - (1) Hemidactylus showing sexual dimorphism
 - (2) Varanus showing mimicry
 - (3) Garden lizard (colotes) showing camouflage
 - (4) Chamaeleon showing protecting colouration
- 172. The figure given below shows three velocity substrate concentration curves for an enzyme reaction. What do the curves a, b, and c depict respectively



- (1) a normal enzyme reaction,
 - b competitive inhibition,
 - c non-competitive inhibition
- (2) a enzyme with an allosteric modulator added, b – normal enzyme activity.
 - c competitive inhibition.
- (3) a enzyme with an allosteric stimulator,
 - b competitive inhibitor added,
 - c normal enzyme reaction.
- (4) a normal enzyme reaction,
 - b non-competitive inhibitor added,
 - c allosteric inhibitor added.
- 173. Pollution from animal excreta and organic waste from kitchen can be most profitably minimized by:
 - (1) storing them in underground storage tanks
 - (2) using them for producing biogas
 - (3) vermiculture
 - (4) using them directly as biofertilizers
- 174. A person who shows unpredictable moods, outbursts of emotion, quarrelsome behaviour and conflicts with others is suffering from:
 - (1) Borderline personality disorder (BPD)
 - (2) Mood disorder
- (3) Addictive disorder
- (4) Schizophrenia
- 175. Genes present in the cytoplasm of eukaryotic cells are found in :
 - (1) mitochondria and inherited via egg cytoplasm
 - (2) lysosomes and peroxisomes

- (3) golgi bodies and smooth endoplasmic reticulum
- (4) plastids and inherited via male gamete
- 176. The type of epithelial cells which line the inner surface of fallopian tubes, bronchioles and small bronchi are known as:
 - (1) squamous epithelium (2) columnar epithelium
 - (3) ciliated epithelium (4) cubical epithelium
- 177. Tadpoles of frog can be made to grow as glant sized tadpoles, if they are:
 - (1) administered antithyroid substance like thiourea
 - (2) administered large amounts of thyroxine
 - (3) reared on a diet rich in egg yolk
 - (4) reared on a diet rich in both egg yolk and glucose
- 178. When children play bare footed in pools of dirty water and flood water, they may suffer from diseases like:
 - (1) Leptospirosis and bilharzia
 - (2) Malaria, amoebic dysentery and leptospirosis
 - (3) Bilharzia, infective hepatitis and diarrhoea
 - (4) Guinea worm infection, elephantiasis and amoebic dysentery
- 179. Which one of the following is an environment related disorder with the correct main cause?
 - Black lung disease (pneumoconiosis) found mainly in workers in stone quarries and crushers
 - (2) Blue baby disease (methaemoglobinaemia) due to heavy use of nitrogenous fertilizers in the area
 - (3) Non-Hodgkin's Lymphoma found mainly in workers involved in manufacture of neem based pesticides
 - (4) Skin cancer mainly in people exposed to benzene and methane
- 180. The figure below shows an angiogram of the coronary blood vessel, Which one of the following statements correctly describes, what is being done?



 It is coronary artery which has a cancerous growth that is being removed (3) Cytokinins

(4) Ethylene

193. What is common between chloroplasts, chromoplasts and leucoplasts?

- (1) Presence of pigments
- (2) possession of thylakoids and grana
- (3) Storage of starch, proteins and lipids
- (4) Ability to multiply by a fission-like process 194. Plants of which one of the following

194. Plants of which one of the following groups of genera are pollinated by the same agency:

- (1) Triticum, Cocos, Magnifera
- (2) Ficus, Kigelia, Casuarina
- (3) Salvia, Morus, Euphorbia
- (4) Bombax, Butea, Bauhinia

195. The Montreal Protocol refers to :

- (1) Persistent Organic Pollutants
- (2) Global warming and climate change
- (3) Substances that deplete the ozone layer
- (4) Biosafety of Genetically Modified Organisms

196. Keystone species deserve protection because these:

- are capable of surviving in harsh environmental conditions
- (2) indicate presence of certain minerals in the soil
- (3) have become rare due to over exploitation
- (4) play an important role in supporting other species
- 197. In India, we find mangoes with different flavours colours, fibre content, sugar content ote: The Information provided here is only

and even shelf-life. The large variation is on account of :

- (1) Species diversity
- (2) Induced mutations
- (3) Genetic diversity (4) Hybridization
- 198. During protein synthesis in an organism, at one point the process comes to a halt. Select the group of the three codons from the following, from which any one of the three could bring about this halt:
 - (1) UUU, UCC, UAU
 - (2) UUC, UUA, UAC
 - (3) UAG, UGA, UAA
 - (4) UUG, UCA, UCG

199. Biosphere Reserves differ from National Parks and Wildlife Sanctuaries because in the former:

- (1) Human beings are not allowed to enter
- (2) People are an integral part of the system
- (3) Plants are paid greater attention than the animals
- (4) Living organisms are brought form all over the world and preserved for posterity

200. Somaclonal variation is seen in :

- (1) Tissue culture grown plants
- (2) Apomicts (3) Polyploids
- (4) Vegetatively propagated plants

for reference .It may vary the Original

Answers with Explanations

1. (3) Tetanus, also known as lockjaw, is a serious but preventable disease that affects the body's muscles and nerves. It typically arises from a skin wound that becomes contaminated by a bacterium called Clostridium tetani, which is often found in soil. Once the bacteria are in the body, they produce a neurotoxin (a protein that acts as a poison to the body's nervous system) known as tetanospasmin that causes muscle spasms. The toxin first affects nerves controlling the muscles near the wound. It can also travel to other parts of the body through the bloodstream and lymph system. As it circulates more widely, the toxin interferes with the normal activity of nerves throughout the body, leading to generalized muscle spasms. Without treatment, tetanus can be fatal

Anti tetanus injection or antitoxin or Equine antitetanus is purified concentrated liquid serum (ATS, tetanus antitoxin, TAT) is a protein fraction isolated from blood of horses hyperimmunized with tetanus anatoxin or toxin.

- 2. (1) Cancer is a class of diseases or disorders characterized by uncontrolled division of cells and the ability of these cells to invade other tissues, either by direct growth into adjacent tissue through invosion or by implantation into distant sites by metastasis. Metastasis is defined as the stage in which cancer cells are transported through the bloodstream or symphatic system. Cancer may affect people at all ages, but risk tends to increase with age, due to the fact that DNA damage becomes more apparent in aging. All these characteristics make the cancer importal until the body in which they resides dies.
- 3. (3) Food chains follow a single path as animals find food. For example: Grass (is eaten by a) GRASS-HOPPER (which is eaten by) a FROG (which is eaten by a) SNAKE (which is eaten by a) HAWK Food webs show how plants and animals are connected to help them all survive. For Example TREES produce acorns which act as food for many MICE and INSECTS. Because there are many MICE the WEASELS. SNAKES AND RACOONS.

have food. The insects in the acorns also attracts SKUNKS, AND OPOSSUMS. BIRDS. SKUNKS, OPPOSUMS, WEASELS and MICE around, HAWKS, KITES, FOXES, and OWLS can find food. They all the corrected.

Thus a network of food chains existing together in an ecosustem is known as a food web.

In the food web kite occupies a specific position because being a carnivorous bird it controls the population of lower carnivores and herbivores.

- 4. (1) In tissue culture, somatic embryos or embryoids are non-zygotic embryo like structure that develop in vitro cultures from somatic cells of any type of tissue, but it is easier to raise them from culture of immature embryos.
- 5. (2) It was generally accepted that the first organisms were heterotrophic (taking food from outside) and fed off a rich prebiotic "soup." Primitive earth was devoid of oxygen so, only those organisms that were able to survive within anaerobic conditions developed. Then after chemo-autotrophs developed that used inorganic sources such as H2S, NH3, CH4 as the principle sources of energy.
- 6. (2) Diarrhoeal disease is an important cause of morbidity and mortality in developing countries, especially among children. In a study of 720 faecal samples from acute diarrhoeal patients various bacterial pathogens were isolated. An account is given of their antibiotic sensitivity pattern. Many of the E. coli, Salmonella and Shigella strains were found to be multidrug resistant. This resistance was transferable and plasmid mediated. A few of the E. coli strains isolated from healthy controls also showed multidrug resistance. Dehydration is common to all types of diarrhoeal diseases. In cases of acute diarrhoea it would be wise to drink more fluids (3-4 litres a day), preferably containing sugar and salts. Ready-mixed rehydration sachets (eg. ORS.Dioralyte, Rehidrat) can be bought from the pharmacist and added to drinking water.
- 7 (2) Global warming is the increase of average world temperatures as a result of what is known as the greenhouse effect. Certain gases in the atmosphere act like glass in a greenhouse, allowing sunlight through to heat the earth's surface but trapping the heat as it radiates back into space. As the greenhouse gases build up in the atmosphere the Earth gets hotter. One of the main greenhouse gases is carbon dioxide (CO2). As trees grow they take in CO2 from the air. When the wood dies the CO2 is returned to the air. Forest clearance and wood burning (such as happens in tropical rain forests) is increasing the latter half of the process, adding to the CO2 in the atmosphere. Deforestation is now out of control. Most people assume that global. warming is caused by burning oil and gas. But in fact between 25 and 30 percent of the greenhouse gases released into the atmosphere each year - 1.6 billion tonnes - is caused by deforestation.

The greenhouse gases selectively transmit the infared waves, trapping some and allowing some to pass through into space. The greenhouse gases absorb these waves and reemits the waves downward, causing the lower atmosphere to warm. Methane stays in the atmosphere for only 10 years, but traps 20 times more heat than carbon dioxide. Fluorocarbons is a general term for any group of synthetic organic compounds that contain fluorine and carbon. Many of these compounds, such as chlorofluorocarbons(CFC's), can be easily converted from gas to liquid or liquid to gas. Because of these properties, CFC's can be used in aerosol cans, refrigerators, and air conditioners. Studies in the 1970's showed that when CFC?s are emitted into the atmosphere, they break down molecules in the Earth's ozone layer (World Book). Since then, the use of CFC?s has significantly decreased and they are banned from production in the United States.

The substitute for CFC's are hydrofluorocarbons (HFC's). HFC's do not harm or breakdown the ozone molecule, but they do trap heat in the atmosphere, making it a greenhouse gas, aiding in global warming. HFC's are used in air conditioners and refrigerators. The way to reduce emissions of this gas is to be sure that in both devices the coolant is recycled and all leaks are properly fixed

Also, before throwing the appliances away, be

sure to recover the coolant in each.

8. (4) Hallucinogens are natural and synthetic (synthesized) substances that, when ingested (taken into the body), significantly alter one's state of consciousness. Hallucinogenic compounds often cause people to see (or think they see) random colors, patterns, events, and objects that do not exist. People sometimes have a a different perception of time and space, hold imaginary conversations, believe they hear music and experience smells, tastes, and other sensations that are not real. So hullucinogens can never be used as analgesics.

Marijuana and hashish, two substances derived from the hemp plant (Cannabis sativa), are also considered natural hallucinogens, although their potency (power) is very low when compared to others. Marijuana (also called grass, pot, tea, weed, or reefer), a green herb from the flower of the hemp plant, is considered a mild hallucinogen. LSD, which is chemically derived from ergot, a parasitic fungus (a fungus that lives in or on a host, deriving benefits from the host while injuring it) that grows on rye and other grains.

(3) Lethal mutation : genetic mutation that kills cells: a genetic mutation with an effect so serious that it often results in the death of the organism concerned. It is a mutant trait that leads to a phenotype incapable of effective reproduction. All types of gene mutations cannot be lethal as most of the mutation are ment for survival and adaptation. Even mutation that is lethal under one condition but not lethal under another condition.

- 10. (1) The Production of T3 and T4 are regulated by thyroid stimulating hormone (T5H), released by the pituitary gland. TSH Production is increased when T3 and T4 levels are too low. The thyroid hormones are released throughout the body to direct the bodies metabolism. They stimulate all cells within the body to work at a better metabolic rate. Without these hormones the bodies cells would not be able to regulate the speed at which they performed chemical actions. Their release will be increased under certain situations such as cold temperatures when a higher metabolism is needed to generate heat.
- 11. (4) In collateral vascular bundles, xylem is situated towards inner side and phloem towards outer side and both are found on same radii. In monocot stem vascular bundle are closed, i.e., cambium is about.
- (3) In Angiosperms, vessel elements in secondary xylem are present for efficient and rapid conduction of water and this conduction is a passive process.
- 13. (1) To increase cell volume, some specialised cells undergo repeated rounds of DNA replication without cell division (endomitosis), forming a giant polytene chromosome. Polytene chromosomes form when multiple rounds of replication produce chromatids that remain synapsed together in a haploid number of chromosomes. They have characteristic light and dark banding patterns which can be used to identify chromosomal rearragements and deletions. Chromosome puffs are diffuse uncoiled regions of the polytene chromosome that are sites of RNA transcription. A Balbiani ring is a large chromosome puff. In addition to increasing the volume of the cells nuclei and causing cell expansion, polytene cells may also have a metabolic advantage as multiple copies of genes permits a high level of gene expression. Polytene chromosomes were originally observed in the larval salivary glands of Drosophila melalanogaster, but are known to occur in secretory tissues of other dipteran insects such as Malphighian tubules of Chironomus and Sciara. They may also occur in protists, plants, mammals, or in cells from other insects.
- 14. (4) Two forms of oxygen are found in the strato-sphere. Molecular oxygen (O₂), which is made up of two atoms of oxygen (O), and ozone (O₃) which, as you can see from its chemical formula, is made up of three oxygen atoms. Ozone is formed when intensive ultra-violet radiation from the Sun breaks down O₂ into two oxygen atoms. These highly reactive oxygen atoms can then react with more O₂ to form O₃.

Global warming and climate changes are mainly due to green-house gases. Ozone holes created continue to grow because of global warming. That's one of the most frustrating things about global warming - its insidious ability to make other environmental problems worse. For example, in addition to having direct effects such as melting polar ice caps and increasing heat waves, global warming also exacerbates seemingly unrelated environmental problems like air pollution, species extinction and holes in the ozone layer.

15. (1) Public concern about global warming mostly focuses on carbon dioxide, the most prevalent greenhouse gas. Methane (CH₄), the major component of natural gas, is second in importance as a greenhouse gas. Methane concentration in the atmosphere has more than doubled during the last 200 years. Its current atmospheric concentration of 1.7 ppm by volume, up from 0.7 ppm in preindustrial times, is much lower than the 345 ppm of carbon dioxide, up from 275 ppm. But one molecule of methane traps approximately 30 times as much heat as does carbon dioxide.

Methane is produced as the terminal step of the anaerobic breakdown of organic matter in wetland rice soils. Methane is exclusively produced by methanogenic bacteria that can metabolize only in the strict absence of free oxygen and at redox potentials of less than -150 mV (which is a property of wetland rice soil)

- (3) At the top is the A horizon, or topsoil, in which humus-unincorporated, often partially decomposed plant residue-is mixed with mineral parti-Technically, humus actually constitutes something called the O or organic horizon, the topmost layer. Examples of humus would be leaves piled on a forest floor, pine straw that covers a bare-dirt area in a yard, or grass residue that has fallen between the blades of grass on a lawn. In each case, the passage of time will make the plant materials one with the soil. Owing to its high organic content, the soil of the A-horizon may be black, or at least much darker than the soil below it. In wetlands of tropical rain forests, the topsoil is shallow and nutrient poor because numerous microorganisms occurring in them quickly break down any litter and the nutrients are recycled immediately to the plants.
- 17. (1) The Gram stain is used to distinguish between gram-positive and gram-negative bacteria, which have distinct and consistent differences in their cel walls. Gram-positive cells may become gram negative through mechanical damage, conversion to protoplasts, or aging, in which autolytic enzymei attack the walls.

In the Gram stain, the cells are first heat fixed and then stained with a basic dye, crystal violet, which is taken up in similar amounts by all bacteria. The slides are then treated with an 1₂-KI mixture (mordant) to fix the stain, washed briefly with 95% alcohol (destained), and finally counterstained with a paler dye of different color (safranin). Gram-posi-

tive organisms retain the initial violet stain, while gram-negative organisms are decolorized by the organic solvent and hence show the pink counterstain. The difference between gram-positive and gram-negative bacteria lies in the ability of the cell wall of the organism to retain the crystal violet.

18. (3) PLANTS absorb light for photosynthesis but as light can itself be dangerous to plants, they need to protect themselves against its damaging effects. Photorespiration is a light induced oxidation of photosynthetic intermediate with the help of oxygen. Photorespiration is stimulated by (i) low CO₂ concentration (ii) high light intensity (iii) high O₂ concentration (iv) high temperature (v) aging of leaf

(Source Internet) Photorespiration results from the oxygenase reaction catalysed by ribulose-1,5bisphosphate carboxylase/oxygenase. In this reaction glycollate-2-phosphate is produced and subsequently metabolized in the photorespiratory pathway to form the Calvin cycle intermediate glycerate-3-phosphate. During this metabolic process, CO2 and NH2 are produced and ATP and reducing equivalents are consumed, thus making photorespiration a wasteful process. However, precisely because of this inefficiency, photorespiration could serve as an energy sink preventing the overreduction of the photosynthetic electron transport chain and photoinhibition, especially under stress conditions that lead to reduced rates of photosynthetic CO2 assimilation. Furthermore, photorespiration provides metabolites for other metabolic processes, e.g. glycine for the synthesis of glutathione, which is also involved in stress protection. In this review, we describe the use of photorespiratory mutants to study the control and regulation of photorespiratory pathways. In addition, we discuss the possible role of photorespiration under stress conditions, such as drought, high salt concentrations and high light intensities encountered by alpine plants.

- 19. (4) Photosynthetically C₄ plants are more efficient than C₃ plants because these have Kranz anatomy (connetive undifferentiated mesophyll around vascular bundles with chloroplast containing bundle sheaths). Bundle sheeth chloroplasts are larger agranal and without PS-II activity and perform CC₃ cycle. (Kalvin cycle)
- 20. (1) Eukaryotic ells have the ability to adopt a variety of shapes and carry out directed movements because they have a network of protein fibres that support the shape of the cell and keep intact the organelles of the cell. This cytoskeleton (network of protein fibres) are made up of actin filaments (microtubulents-stress fibres), microtubules and intermediate filaments.
- (2) Both the assertion and reason are independently true but the reason does not satisfies the

assertion. Starch is used in iodometric titration as an indicator because starch gives blue colour with free iodine. This blue colour disappears when free iodine is completely changed to iodide ion.

 (1) As the bond order (B.O) increases bond length decreases and vice versa.

B. O of
$$N_2 = \frac{1}{2}$$

(No. of bonding electron - No. of antibonding electron)

$$=\frac{1}{2}(6-0)=3$$

B. O of
$$O_2 = \frac{1}{2}$$

(No. of bonding - No. of antibonding electron)

$$=\frac{1}{2}(6-2)=2$$


 $B.\ O\ of\ N_2\ is\ 3$ and $B.O\ of\ O_2\ is\ 2$, therefore, bond length of $N_2\ is\ less\ than\ O_2$

- 23. (1) Optical isomerism only can be shown by the compound which are dis-symmetrical or assymmetrical. The compound [CO (NO₂)₃ (NH₃)₃] is symmetrical because it has plane of symmetry therefore, it will not show optical isomerism.
- 24. (2) Down the series E° value becomes more negative. In the electrochemical series chromium present below Mn, that is why Mn has more positive E° value than chromium.

Third ionisation potential of Mn is larger tan third 1P of Cr because the electronic configuration of Mn^{2+} is $3d^5 4s^6$ and Cr^{2+} is $3d^4 4s^6$. Mn^{2+} has half filled $3d^5$ configuration which is more stable than $3d^4$ (less than half filled) configuration of Cr^{2+} therefore, it is difficult to remove the next electron (3^{nd} electron) from Mn^{2+} ($3d^5$ configuration) as compare to $3d^4$ of Cr^{2+} hence, third IP of Mn is more than third IP of Cr.

- 25. (3) It is true that K₂ Cr₂ O₇ (potassium dichromate) is used as a primary standard in volumetric analysis because it is not hydroscopic in nature and therefore, it does not gain weight by absorbing moisture which leads to purity in state and hence can be measured accurately. But K₂ Cr₂ O₇ is less soluble in water.
- 26. (3) Silicons are polymeric organosilicon derivatives containing Si O Si links. Silicons are inert and strongly water repellent. The tendency for such inertness towards acid, bases, alkalis and water repellent is due to the presence of larger (bulky) organic groups which surrounds the silicone chain.
- 27. (1) According to transition state theory, for the formation of an activated complex one of the vibrational degrees of freedom is converted in to a transitional degree of freedom. It is because the energy of activated complex is higher than the energy of reactant molecules

- 28. (1) At room temperature, fusion of ice is a spontaneous process, because the process again proceeds in that direction where randomness increases. Liquid state (water) is more random (i. e., larger entropy) than solid state (ice). Hence, the melting of ice into water is spontaneous process.
- (4) The assertion is wrong because Sb₂ S₃ is soluble in yellow armmonium sulphide.
 Sb₂ S₃ + 3 (NH₄)₂S —— 2 (NH₄)₃ Sb S₃



The above figure shown sheet like hexagonal structure of graphite in which each carbon atom is sp^2 hybridized. For tetragonal system $a = b \neq c$, $\alpha = \beta = \gamma = 90^\circ$. But for a hexagonal system $a = b \neq c$, $\alpha = \beta = 90^\circ$, $\gamma = 120^\circ$.

31. (2) Both the statements are true independently but reason does not satisfies the assertion. The opposite potential if larger than cell potential than the direction of flow of current will be reversed and it will reverse the cell reaction also, it means where oxidation was vice versa. Now zinc ion are converted to zinc (s) (reduction) at one electrode and Cu is converted in to Cu²⁺ ion (oxidation).

 Zn^{2*} (aq) + Cu (s) \longrightarrow Cu^{2*} (aq) + Zn (s)

- 32. (3) It is true that Fe³⁺ ion can be used for coagulation of As₂S₃ sols. It is due to the fact that when different or oppositely charged sols are mixed together in nearly equal proportions both the sols may be precipitated totally or partly. When As₂ S₃ (negatively charged) is added to Fe(OH)₃ (positively charged) sols, both the sols get precipitated simultaneously if such addition of sols are volumetrically equal or nearly equal.
- 33. (3) Due to osmosis pure water will enter into the blood cells, which led to increase in internal pressure therefore cell swells. If we see the concentration of salt in the blood cell, it will decrease after osmosis.
- 34. (3) Ethyl alcohol is volatile and comes out with the Breath of drunker. When these small vapour of ethyl alcohol is treated with or get contact with acidic K₂ Cr₂ O₇ solution, the colour of the solution changes from orange to green.

The reaction involved in above alcohol drunk test

$$\begin{array}{c} \text{CH}_3 \text{ CH}_2 - \text{OH} \xrightarrow[O]{\text{II}_3 \text{ Cr}_3} \text{O, 14,SO}_4 \\ \text{(BMyl abrolic)} & \text{(ORANGE COLOUR)} \\ \\ \xrightarrow{\text{II}_3 \text{ Cr}_3} \text{O, 14,SO}_4 \\ \text{(ORANGE COLOUR)} & \text{CH}_3 \text{COOH}_4 \\ \text{(ORANGE COLOUR)} \end{array}$$

Here $K_2Cr_2O_7$ acts as cooldising agent in the presence of H_2SO_4 . $K_2Cr_2O_7$ oxidises CH_5 CH_2 OH and isself get reduced. The reduced part becomes $Cr_2(SO_4)_3$, chromic sulphate which is green in colour.

$$K_2Cr_2O_7 + 4 H_2SO_4 \longrightarrow K_2SO_4 + Cr_2 (SO_4)_5$$
(Change)
$$+ 4 H_2O + 3 [O]$$

35. (3) It is true that anilinium chloride is more acidic than ammonium chloride. But anilinium ion is less resonance stablised than aniline, because anilinium ion does not contain lone pair of electrons, therefore resonance is not possible.


Stronger acid has weaker conjugate base and vice versa, on the basis if above fact, anilinium ion is stronger acid because aniline is weaker conjugate base.

- 36. (3) Diastereomers Stereoisomers which are not mirror images of each other are called diastereomers. These compounds have different physical property.
- 37. (1) Both the statement is correct and reason explains the assertion. The nucleophilic substitution can occur when strong electron-withdrawing groups are ortho or para to halogen atom.

The mechanism that operates in these reactions is an addition - elimination mechanism involving the formation of a carbanion with delocalised electrons called a Meisenheimer complex.

(Meisenheimer Complex)

Carbanion is stabilized by electron - withdrawing groups (- NO₂) in the positions ortho and para to the halogen atom



Especially stable (-we charges are both an oxygen atom)

38. (4) 1, 3-Butadiene is not a monomer of natural rubber but it is isoprene i. e., 2-methyl-1, 5-butadiene. It is a general rule that unsaturated monomer when polymerises it gives poly addition product. Such addition in the case of natural rubber is through cationic addition polymerization. The reaction involved in cationic addition is

n
$$CH_2 = \overset{\circ}{C} - CH = CH_2 \xrightarrow{Polymerisation}$$
 CH_3
 $\leftarrow CH_2 - \overset{\circ}{C} = CH - CH_2 + \overset{\circ}{n}$

$$CH_3 - CH = CH - CH_3 \xrightarrow{HSC} CH_3 - CH - CH_2 - CH_3 \xrightarrow{BC} CH_3 - CH_2 - CH_3 \xrightarrow{BC} CH_3 - CH_3 - CH_3 - CH_3 \xrightarrow{BC} CH_3 - CH_3$$

Above reaction gives only one product i. e., 2-bromobutene

- 40. (1) Addition of non-volatile solute in to any volatile solvent reduces the vapour pressure. Such reduce in vapour pressure will be in equilibrium with solid phase at lower pressure and hence at a lower temperature. Such addition of non-volatile solute in to solvent results in to depression of freezing point of the solvent.
- 41. (3) For a wave to propagate in sky, it has to satisfy the principle of sky wave propagation. According to the principle, if the frequencies of the waves are less than the critical frequency then the waves would be reflected back since the ionosphere has refractive index less than free space. Thus the assertion is wrong. The statement given in reason is correct but it is not true for the assertion
- (2) For heavy nuclei (A>100), the binding energy per nuclear decreases due to coulomb repulsion between the protons inside the nucleus
- 43. (3) The assertion is true because the common base configuration of npn transistor is used for voltage amplification with the current amplification being very small.


The reason is incomplete and is thus wrong because the collector is reverse biased for voltage amplification which has not been mentioned.

 (2) The entropy of an isolated system increases in accordance with the second law of thermodynamics.

- (1) The phenomenon of nuclear magnetic resonance is used in Magnetic Resonance Imaging, thus both assertion and reason are correct.
- 46. (1) The judo fighter initially has to try to bend his opponent and then rotate him around his hip because, then the opponents moment of inertia would be reduced and due to the any torque given by a force would be resisted but not the torque by rotation.
- 47. (4) The root mean 'square and the most probable speeds are given by v_{max} = √3 RT/M

and
$$v_{propatable} = \sqrt{\frac{2 RT}{M}}$$

and the maxwellian distribution of velocity is given by which is not symmetrical.



- 48. (3) The ball bearing is used to reduce the friction
- (1) Gratings produced diffraction patterns and they cannot distinguish different wavelengths as their spacing is not of the same order.
- 50. (4)
- (2) A man inside a freely falling lift falls with zero relative acceleration

The reason is correct but is not a correct explanation

52. (3) The reason is wrong as the work function is dependent only on the photoelectric metal.

53. (1) 54. (1)
55. (3) As the skin loses water due to perspiration, it enhances its emissivity.

56. (2) The reason is not the correct explanations

57.(1) 58.(2) 59.(1) 60.(1)

61. (2) us = vn (n+2)

where μ s = spin only magnetic moment n = number of unpaired electrons

if number of unpaired electrons are same µs will be same. In Cr²⁺ and Fe²⁺ the number of unpaired electron is 4 therefore µs is same. Co²⁺ has 3 unpaired electrons and Mn²⁺ has 5 unpaired electrons.

62. (4) Haemoglobin (Haem - ion; globin - globular protein) contain iron as central atom in prohyprin ring. Cytochromes are widely distributed respiratory catalyst concentrated in electron transport chain of living cells. Cytochromes are haemoproteins differing in prophyrin groups. It also contains iron in its prophyrin ring.

63. (1) Na₂ B₄ O₇ (Borax) when dissolves in water and gives its alkaline solution. Because of this reason borax is used as water softening and cleansing agent

Sodium hydroxide cleans and softens the water.

64. (3) Sulfur (Sulphur) exist in several allotropic forms i. e., α, β and γ sulphur. All the three forms contains Puckered S₈ ring with crown conformations. But Engel's sulphur (ε- sulphur) contains S₅ rings. Some other form of sulphur (allotropes) have been synthesized are

65. (1) Sodium bicarbonate and magnesium hydroxide are used to prepare antacid or anti acid medical preparations.

$$NaHCO_3 + HCI \longrightarrow NaCI + H_2O + CO_2$$

 $Mg(OH)_2 + 2 HCI \longrightarrow MgCI_2 + 2 H_2O$


Both the compounds reacts with HCl and neutralizes the acids and increase the pH of stomach.


- 66. (2) The glass is an amorphous substance which is clear (watery) solid. It can some time be coloured after mixing transition element (especially its ions). On mixing Co²⁺ ion in the glass gives deep blue colour.
- 67. (1) The central atom in cisplatin is Pt and the ligands are Cl and NH₃. The IUPAC name of cisplatin is cis-diammine dichloroplatinum (II).



It is used as anticancer drug. The structure of cisplatin is

68. (3) [Co(C2O4h (NH3)2]






trans-isomer


- 69. (3) In Haber Bosch process ammonia gas is prepared by reacting H₂ and N₂ in the presence of finely divided iron (catalyst) and molybdenum (promoter).
- 70. (1) XeF₄ involves sp³ d² hybridisation and its shape square planar instead of octahedral due to presence of two lone pair of electron of Xe atom. Here two lone pair of electron will occupies the polar position (according to VSEPR theory)



SF₄ molecule involved sp³ d hybridisation and due to presence of one lone pair of electron the distorted geometry of see-saw like structure instead of trigonal bipyramidal.




[NiCl₄]²⁻ ion involved sp³ hybridisation as follows



sp³ hybridisation

Due to sp³ hybridisation its shape is tetrahedral. [pd Cl₄]²⁻ ion involved d sp² hybridisation as follows



Due to the d sp² hybridization [pd Cl₄]²⁻ ion shows square planar geometry.

- 71. (2) Tincture iodine is a solution of l2 in aqueous KI.
- 72. (3) [Ag (C ^a/_σ N)₂]⁻ has 4 π bond which is clear from the diagram.

- 73. (1) The compound phosphorus pentachloride (PCI₅) behaves as molecule in gaseous phase. In solid state it acts as ionic solid as
 - [PCI4]* [PCI6]*
- 74. (1) Potassium iodate and potassium iodide salts are used to prepare iodide salt when its traces are mixed with large amount of NaCl
- (1) U₃ O₈ is the compound used in enrichment of Uranium for power plant. These U₃ O₈ contains Uranium - 235

76. (1) K.E =
$$\frac{1}{2}$$
 m V^2

(where m = mass, V = velocity)

$$\Rightarrow 0.5 = \frac{1}{2} \times 1 \times V^2$$

⇒ V = 1 m/s

de Broglie wavelength,

$$\lambda = \frac{h}{m V} = \frac{6.626 \times 10^{-34} \text{ Js}}{1 \text{ Kg} \times 1 \text{ ms}^{-1}}$$

$$\lambda = 6.626 \times 10^{-34} \text{ m}$$

- 77. (2) If the compressibility factor is Z then dominance of strong repulsive force depends on Z > 1.
 If Z < 1 it is due to attractive force.</p>
- 78. (3) 40 ml of 0.1 M NH₃ solution = 40 × 0.1
 - ⇒ 4 milli equivalent ammonia solution

 $20 \text{ ml of } 0.1 \text{ M HCl} = 20 \times 0.1$

2 milli equivalent of HCl

For the reaction

$$NH_4OH + HCI \longrightarrow NH_4CI + H_2O$$

Initially 4 2 0
After reaction 4 – 2 0 2
= 2

$$pOH = pKb + log \frac{[NH_4CI]}{[NH_4OH]} = 4.74 + log \frac{2}{2}$$

79. (4) Total change in entropy,

For spontaneous process ΔS_{cont} must be positive.

$$\Delta S_{\text{res}} > 0$$

- Total entropy change is always positive if process is spontaneous.
- 80. (1) The Ca²⁺ ions are arranged in ccp arrangement. In this arrangement Ca²⁺ ions are present at all corners and at the centres of each face of the cube. The fluoride ion (F) occupy all the tetrahedral site. There are two tetrahedral sites per atom in a closed packed lattice. This means that there are two tetrahedral sites every Ca²⁺ ion. Since F ions occupy all the tetrahedral sites, there will be two Fions. Thus the formula of the compound is CaF₂.
- 81. (2) MnO₄ + 2 H₂O + 3 ē ---→ MnO₂ + 4 OH⁻ It is clear from the above equation that for the reduction of one mole of MnO₄ ion to MnO₂ 3 electron is required. It is because the oxidation state of Mn in MnO₄ ion is + 7 and the oxidation state of Mn in MnO₂ is + 4.
- 82. (3) For the reaction

$$x_1 A + x_2 B \longrightarrow y_1 C + y_2 D$$

Rate expression for the above reaction is

$$-\frac{1}{x_1} \frac{d[A]}{dt} = -\frac{1}{x_2} \frac{d[B]}{dt} = \frac{1}{y_1} \frac{d[C]}{dt}$$
$$= \frac{1}{y_2} \frac{d[D]}{dt}$$

Now, for the reaction

Rate expression is

$$-\frac{1}{2}\frac{d \left[N_{2} O_{5}\right]}{dt} = \frac{1}{4}\frac{d \left[NO_{2}\right]}{dt} = \frac{d \left[O_{2}\right]}{dt}$$

83. (1) For the reaction

For a phase change $\Delta G = 0$

Because $\Delta G^{\circ} = RT \ln K$

[\(\)G^0 = standard free energy change, K=equilibrium constant]

If a substance is in equilibrium between two phases at constant temperature and pressure the free energy change is zero.

84. (3) For water
$$K_f = \frac{\Delta T_f \times W \times m}{1000 \times \omega}$$

(Where W=weight of water, ω=wt. of cane sugar, m=molecular wt. of cane sugar)

$$K_t = \frac{2.15 \times 100 \times 342}{1000 \times 5} = 14.7$$

For 5% C₆ H₁₂ O₆

$$\Delta T_f = \frac{K_f \times 1000 \times \omega'}{W \times m'}$$

(where \omega' = wt. of C6H12O6, m' = molecular wt. of C, H12 O6)

$$\Delta T_f = \frac{14.7 \times 1000 \times 5}{100 \times 180} = 4.08$$

Now, freezing point of C₆ H₁₂ O₆ solution = 273 - 4.08 = 269.07 K

85. (1) The order of energy gap is

Egypternond) > Egyptern) > Egypternond As the energy gap (Eg) increases conductivity de-CTRRSS.

86. (2) The change in enthalpy AH = AU + An RT

(where $\Delta n = no$, of gaseous product - no. of gaseous molecule)

-92.38 × 1000 = ∆U - 2 × 8.314 × 298

(Reactant is more in number than product so negative sign is there)

 $\Delta U = -87424 J = -87.424 kJ$


87. (3) For the electrolysis of the salt

At cathode,
$$2 H^+ + 2 e^- \rightarrow H_2 \uparrow$$

At anode, 2 Br - 2 e → Br₂ ↑

In solution, Na* + OH⁻ → NaOH

88. (3) According to corn rule, L-serine is



L . Serine

2-amino-3-hydroxypropanoic acid. It the priority groups i.e., - NH2, - COOH, - CH₂OH, - H are arranged clockwise it is said to be a L - form.

89. (2)

92. (4)

- NHAc > - CH₂ > X


93. (2) It two o and p - directing group is attached to an aromatic ring the directing power of attached group is in the order of 0" > - NH₂ > - NR₂ > - OH > - O - CH₃ >

Compound (I) will be preferred of less steric effect.

95.(1) Thymine is 5-methyluracil. Its structure is

Thymine. (A nitrogenous base in nucleic acid.

- (4) Lysin is an aminoacid, it is soluble in law pH or in acidic medium but it is less soluble in isoelectric pH i. e., basic pH or high pH.
- (2) Methyl α D glucoside and methyl β D-glucoside are anomers because they differs only



(2) Boiling point increases if size of alkyl group increase and boiling point decreases if branching increases.

So the order of boiling point.

> CH,CH(CH,)CH,CI > (CH,),C CI

 (1) The increasing order of reactivity of halides for S_N1 reaction is

$$CH_3 - CH_2 - x \le (CH_3)_2CH - X$$

 $\le CH_2 = CH - CH_2 - X \le PhCH_2 - X$

In this reaction halogen atom is replaced by - OH group (nucleophile).

- (3) Light bends since the speed of light is different in different media.
- 102. (4) The acceleration due to gravity is independent of the mass
- 103. (2) The moment of inertia of the rod about its centre and perpendicular to its axis is ¹/₁₂ ML² and

the moment of inertia about one end = $\frac{ML^2}{3}$, so the moment of inertia about 0 is

$$V1$$
 $m/2$
 $m/2$
 $m/2$
 $M \frac{(l/2)^2}{3} + \frac{M (l/2)^2}{3} = \frac{ML^2}{12}$

104. (3) Given λ = 100 m, v = 25 m/sec

then the frequency = $\frac{25 \text{ m/sec}}{100 \text{ m}} = \frac{1}{4} \text{sec}^{-1}$

So the time of bounce = 4 sec

105. (3) The atm a spheric pressure = 760 mm of Hg The pressure in the lungs = 750 mm of Hg so the difference in pressure = 10 mm

= 1 cm of Hg and the density = 13.6 g/cm^3 so we have $1 \times 13.6 \times \text{g} = l \times 1 \times \text{g}$

o /= 13.6 cm

Thus water can be drawn from a depth of 13.6 cm. 106. (3) The surface charge densities on the sheets are given by $\sigma = 26.4 \times 10^{-12} \, \text{c/m}^2$ so the electric field between the sheets is

$$E = \frac{\sigma}{\epsilon_0} = \frac{26.4 \times 10^{-12}}{8.85 \times 10^{-12}} \approx 3 \text{ N/C}$$

- 107. (2) The dimension of magnetic moment is [L² A]
- 108. (1) Given u = -8 cm. f = 10 cm

so using
$$\frac{1}{v} - \frac{1}{u} = \frac{1}{f}$$

we get $\frac{1}{v} = \frac{1}{u} + \frac{1}{f}$
 $= \frac{-1}{8} + \frac{1}{10} = \frac{-10 + 8}{80} = \frac{-2}{80}$

hence the magnification produced by the lens is

$$m = \frac{v}{u} = \frac{-40}{-8} = 5$$

109. (2) Given \(\lambda_{\text{min}} = 10^{11} \text{ m.}\)

So
$$V_{max} = \frac{hC}{e \lambda_{min}} = \frac{1.242 \times 10^{-6}}{10^{-11}}$$

so $V_{max} = 124.2 \text{ kV}$

Which is the accelerating voltage.

 (2) The electrons are ejected if the incident light has certain minimum frequency

111. (2) Given aug = 1.5

then the focal length of lens by 'f'

so
$$\frac{1}{f} = (a_{\mu g} - 1) \left(\frac{1}{R_1} - \frac{1}{R_2} \right)$$

 $\frac{1}{f} = 0.5 \left(\frac{1}{R_1} - \frac{1}{R_2} \right)$...(i)

when the lens is immersed in a liquid of refraction index 1.25, then

$$\frac{1}{f'} = 0.25 \left(\frac{1}{R_1} - \frac{1}{R_2} \right)$$

sof' = 2.5f

112. (1) The voltage on the clouds = 4 × 10⁶ volts Time of strike = 100 m sec

Charge delivered = 4 coulomb

So the power lightening = $\frac{\text{Work done}}{\text{Time}}$ = $\frac{4 \times (4 \times 10^6 - 0)}{100 \times 10^{-3}}$ = $\frac{16 \times 106}{100 \times 10^{-3}}$ = 160 mega

113. (3) Between P and R, a series combination of two capacitances in parallel with a series combination of three capacitance. Thus equivalent capacitance between P and R is

 $C_{p_1} = \frac{C}{3} + \frac{C}{2} = \frac{5C}{6}$

Similarly between P and Q, a parallel combination of one capacitance with a series combination of four capacitance

Thus
$$C_{PQ} = C + \frac{C}{4} = \frac{5 C}{4}$$

So $\frac{C_{PR}}{C_{PQ}} = \frac{4}{6} = \frac{2}{3}$

114. (1) For a circular wave, the amplitude of the wave varies as A \(\alpha \) r^{-1/2}

115. (4) In the inelastic collision between two spherical rigid bodies, the linear momentum is conserved but not the kinetic energy 116. (2) When a p-n diode is reverse biased, then the depletion region of the p-n diode is increased

117. (1) When the multiplication factor has a value less than 1, then the operation of the nuclear reactor is said to be critical

118. (2)

119. (1) The output = product of inputs

120. (3) To move a load with constant velocity, first the force is used to overcome the stastic friction and then the body keeps on moving with constant velocity even when the force has been decreased.

121. (2) The black object will glow the brightest 122. (2) It it is assumed the balloons have same volume, so using PV = nRT, and given is P, T are same, so h' is also same

123. (4) As the battery wears out the filament shows red colour though if there is no significant change in voltage, colour remains the same but its intensity decreases.

124. (1) The lines of forces move out from a positive charge and go into a negative charge.

125. (1) The magnetic field due to a circular loop at its centre is $B' = \frac{\mu_0}{4\pi} \frac{2\pi I_C}{R}$ and that due to a straight

line at the centre of the loop is B" = $\frac{\mu_0 \times 2 I_C}{4 \pi H}$

so
$$B = B' - B'' = 0$$

so $H = I_c R / \pi I_c$

126. (1) The moment of the tension about the is maximum in A and it counter balances the moment of mg acting from the centre of the rod.

127. (2) The emission of an α particle is followed by

(i) Decrease in mass number = 238 - 4 = 234 (ii) Decrease in atomic number = 92 - 2 = 90 So 234 Th is emitted.

128. (3) Given the amount of C^{14} remaining is $\frac{1}{2}$ of origin after 5730 years. So the have $\frac{1}{16}$ th of origin

nal value, it takes
$$\left(\frac{1}{2}\right)^4$$

⇒ 4 × 5730 = 22920 years.

129. (3) By conservation of mass number and charge no. ²³Na and ¹H is the right answer.

130. (1) For beats of 5Hz, 440 Hz tuning fork is used for beats of 8Hz, 432 Hz tuning fork is used so 445 Hz is the correct small.

131. (2) (i) When the ring enters the magnetic field the ernf is induced and current flows in one direction.

(ii) When the ring is inside the filled, there is no change in flux.

(iii) When the ring gas out, the current is in opposite direction to that in (i)

- 132. (4) For α and β, the wavelength will be de Broglie wavelength and for γ waves, the wavelength would be that of electromagnetic waves.
- 133. (2) The voltage gain in dB = 20 log₁₀ A, = 20 log₁₀ (1000) = 20 × 3 = 60 B
- 134. (3) When water is converted to ice, entropy decreases.
- 135. (2) In cold bath, length of x decreases more, so it moves towards left.
- 136. (3) For a wave propagating in a medium the frequency is independent
- 137. (2) If an object reflects the colour of light incident on it, it will appear with that colour but if object absorbs the colour of light, it will appear to be black. Since, the given wavelength does not belong to green, so it will be absorbed by the leaf and hence, it will appear to be black.
- 138. The voltage across its ends = 2 v so the pet. drop across R = 6 - 2 = 4v Current flowing = 10 mA

So R =
$$\frac{4}{10 \times 10^{-3}}$$
 = 400 Ω

- 139. (1) The potential difference between base and emitter is 1V
- 140. (4) For the demodulation to be good, RC > $\frac{1}{6}$.
- 141. (d) CPR: cardiopulmonary resuscitation, a combination of rescue breathing (mouth-to-mouth resuscitation) and chest compressions. If a child isn't breathing or circulating blood adequately, CPR can restore circulation of oxygen-rich blood to the brain. Without oxygen, permanent brain damage or death can occur in less than 8 minutes.
- 142. (4) Alzheimer's (AHLZ-high-merz) disease is a progressive brain disorder that gradually destroys a person's memory and ability to learn, reason, make judgments, communicate and carry out daily activities. It is common in elderly people.
- 143. (1) The Govt. of India after having recognized the need for systematic Research and Development of Ayurveda, Yoga & Naturopathy, Unani, Siddha and Homoeopathy in the country, established the "Central Council for Research in Ayurveda, Yoga & Naturopathy, Unani, Siddha and Homoeopathy" in 1969 as an autonomous organization under the Ministry of Health and Family Welfare, 61-65, Institutional Area, Janakpuri, Phankha Road, New Delhi-110058
- 144. (4) The term pneumonia refers to any infection of the lung. The "pneumonia vaccine" is given to prewent one specific type of pneumonia-the pneumonia caused by the Pneumococcus (Streptococcus pneumoniae) bacterium. Vaccination for pneumonia is not covered in the immunization schedule so far in India.

DTaP is an acronym for the combined diphtheria, tetanus, and acellular pertussis vaccine. The "a" de-

notes the vaccine's acellular pertussis components, distinguishing it from whole-cell, inactivated DTP (aka DTwP) vaccine. The acellular vaccine uses antigenic fragments of the pertussis pathogen to induce immunity

MMR/Measles Vaccine is given for protection against measles, the typical fever with rash of childhood

145. (1) It is an old wives' tale. Really. Try it with 100 cucumbers and not on another 100 if you don't believe me. Some cucumbers can be bitter, especially at the ends, and you might want to cut off the ends because of this. In particular, cut off the blossom end of fresh cucumbers, as the blossoms contain enzymes that are bitter and others that cause excessive softening of pickles.

You can draw out the bitterness in the rest of the cucumber by slicing, salting, and draining it before adding to a salad. This works on the principle of osmoss. The principle of bitterness spontaneously leaving the cucumber because you rubbed the end... well, it does no harm, so do it if you want.

146.(1) Rustom Jal Vakil returned to India in 1938 after earning his medical degree from the University of London and focused on the treatment of heart ailments at a time when cardiology was not a distinct subspecialty in India. In 1949, after years of scrupulous collation and analysis of data, he published a watershed paper on the antihypertensive properties of Rauwolfia serpentina and effected a paradigm shift in the management of hypertension. Rauwolfia was the world's 1st successful bloodpressure-lowering agent, and its acceptance encouraged research scientists to pursue the development of other hypotensive drugs. Reserpine was isolated in 1952 from the dried root of Rauwolfia serpentina (Indian snakeroot), and introduced in 1954, two years after chlorpromazine. Reserpine almost irreversibly blocks the uptake (and storage) of noradrenaline and dopamine into synaptic vesicles by inhibiting the Vesicular Monoamine Transporters (VMAT).

Paul Ehrlich (March 14, 1854 – August 20, 1915) was a German scientist who won the 1908 Nobel Prize in Physiology or Medicine. He is noted for his work in hematology, immunology, and chemotherapy. Ehrlich predicted autoimmunity and called it "horror autotoxicus". He coined the term "chemotherapy". The idea of a "magic bullet" is also his, and he is credited with the first empirical observation of the blood-brain barrier.

147. (2) Biometry or biometrics is simply defined as automatically recognizing a person using distinguishing traits. Biometric identification uses some unique physiological or behavioural characteristic, not shared by any other individual, to positively identify an individual. . 148. (2) Each day after the winter solstice, which occurs on December 21st, the Sun's path becomes a little higher in the southern sky. The Sun also begins to rise closer to the east and set closer to the west until we reach the day when it rises exactly east and sets exactly west. This day is called the equinox. In the spring we have the Spring Equinox about March 21st. There is also a Fall Equinox on September 21st

149. (1) In late 1895, a German physicist, W.C. Roentgen while working with a cathode ray tube in his laboratory discovered X-rays.

150. (2)

151.(3) Directly Observed Therapy Shortcourse (DOTS) is the internationally-recommended TB control strategy and is composed of five distinct elements: political commitment; microscopy services; drug supplies; surveillance and monitoring systems and use of highly efficacious regimens; and direct observation of treatment.

152. (4) 153. (1) 154. (2) 155. (3) 156.(1) 157. (1) 158. (1) 159. (2) 160. (2)

161. (2) The autonomic nervous system (ANS) is the part of the nervous system of the higher life forms that is not consciously controlled. It is commonly divided into two usually antagonistic subsystems: the sympathetic and parasympathetic nervous system, and involves the homeostasis of organs and physiological functions.

A third and less commonly considered part of the autonomic nervous system is the enteric nervous system, which controls the digestive organs, and is, for the most part, independent of central nervous system (CNS) input.

In general, the parasympathetic nervous system is involved with digestion and energy conservation, while the sympathetic nervous system is involved with energy expenditure and the 'fight or flight' re-

Here is a summary of some of the effects of sympathetic and parasympathetic stimulation. Notice that effects are generally in opposition to each other.

The Autonomie Me

Structure	Sympathetic Stimulation	Parasympathetic Stimulation
iris (eye muscle)	Pupil dilation	Pupil constriction
Salivary Glands	Saliva production reduced	Saliva production increased
Oral/Nesal Mucosa	Mucus production reduced	Mucus production increased
Heart	Heart rate and force increased	Heart rate and force decreased
Lung	Bronchial muscle relaxed	Bronchial muscle contracted
Stomach	Peristalsis reduced	Gastric juice secreted; motility increased

Small Intestine	Motility reduced	Digestion increased	
Large Intestine	Motility reduced	Secretions and motility increased	
Kidney	Decreased urine secretion	Increased urine secretion	
Liver	Increased conversion of glycogen to glucose		
Adrenal medulla	Norepinephrine and epinephrine secreted		
Bladder	Wall relaxed	Wall contracted	
DESCURIT	Sphincter closed	Sphincter relaxed	

It should be noted that the autonomic nervous systern is always working. It is NOT only active during "fight or flight" or "rest and digest" situations. Rather, the autonomic nervous system acts to maintain normal internal functions and works with the somatic nervous sustem.

162. (2) The Duckbill Platypus' are mammals because the females produce milk and nurse their young; however they are different as the Duckbill platypus, strangely lays eggs. This makes the Duckbill Platypus one of only three Monotremata that walk this earth. Originating from the word monotreme, which means a "single whole or opening", Monotremata are mammals that lay eggs. The Duckbill platypus is a mammal that lays eggs.

(1) Despite their name (which means "thousand legs"), millipedes do not have 1,000 legs - they have from 47 to 197 pairs of legs, depending on the species. Millipedes are invertebrates; they have a hard exoskeleton and many jointed legs. The nervous system comprises a three-part brain, with a nerve cord running the body's length, with bundles of coordination nerves. Classification: Kingdom Animalia (animals), Phylum Arthropoda (jointed legs and an exoskeleton), Superclass Myriapoda (*manyfooted" with a 2-segmented body - millipedes. centipedes, etc.), Class Diplopoda (millipedes). About 7,500 species of millipedes have been described.

(3) Lepisma saccharina

Phylum: Arthropoda Class: Insecta

> Order: Thysanura Family: Lepismatidae Genus: Lepisma

Species: L. saccharina

Lepisma saccharina (commonly called the fishmoth, urban silverfish or just the silverfish) is a small, wingless insect. Its common name derives from the animal's silvery grey colour, while the scientific name indicates the silverfish's diet of carbohydrates such as sugar or starches. It belongs to the basal insect order Thysanura, and the species is estimated to have existed for 300 million years.

(4) Sea anemone are Diploblastic: having a body made of two cellular layers only (as opposed to the majority of animals which are triploblastic), the exterior ectoderm and the interior endoderm. These two layers are separated by the largety non-cellular jelly-like mesogloce.

Classification: Kingdom Animalia (animals), Phytum Coelenterata (corals, jellyfish, sea anemones, hydroids), Class Anthozoa meaning "flower-like animals" (corals and sea anemones), Order Actiniaria

nes), Order Actiniana

(4) muscular diaphragm and milk producing glands.

(1) monotremes are oviparous

- (2) Elephants are one of the exceptions to the rule that mammals have extra-abdominal testes; so are whales and perhaps other marine mammals.
- (3) All mammals except the monotremes, the edentates, the pangolins, and the cetaceans have up to four distinct types of teeth, with a maximum number for each. These are the incisor (cutting), the canine, the premoler, and the molar (grinding). Mammals that have distinct types of teeth are historodost, others are homodorst. Mouth with diphyodont teeth (milk, or deciduous, teeth replaced by a permanent set of teeth); teeth in most spp. are heterodont (teeth with various structures and functions; lower jaw a single enlarged bone (dentary).

General Characteristics of Mammals

- Body covered with hair, but reduced in some spp..
- Integument with sweat, scent, sebaceous, and mammary glands.
- 3. Skeletal features include:
 - a. skull with
 - I. bony palate
 - middle ear with three ossicles (malleus, incus, and stapes)
 - b. Seven cervical vertebrae
 - c. Fused pelvic bones.
- 5. Movable eyelids and fleshy external ears
- Four limbs (reduced or absent in some spp.) adapted for many forms of locomotion.
- Circulatory system of 4-chambered heart, nonnucleated RBC's, and biconcave RBC's.
- Respiratory system of lungs with alveoli, and voice box (larynx); a secondary bony palate separates air and food passages; muscular dia-

- phragm separates thoracic and abdominal cavi-
- Excretory system of metanephric kidneys and ureters that usually open into a bladder.
- Brain highly developed; 12 pairs of cranial nerves.
- Strictly endothermic homeotherms.
- Cloaca present only in monotremes (egg-laying mammals, such as duck-billed platypus).
- Separate sexes; specialized reproductive and copulatory organs.
- Internal fertilization; egg develops in a uterus with placental attachment (except in monotremes); fetal membranes (amnion, chorion, allantois); sex determination by heterogametic males (females are isogametic).
- Young are nourished by milk from mammary clands.
- 164. (1) Parietal cells (also called oxyntic cells) are cells located in the stomach epithelium.

Parietal cells contain an extensive secretory network (called canaliculi) from which the HCl is secreted by active transport into the stomach. The enzyme hydrogen potassium ATPase (H*/K* ATPase) is unique to the parietal cells and transports the H* against a concentration gradient of about 3 million to 1.

- (2) Alpha cells are located in the selets of Langerhans in the pancreas; they produce the hormone glucagon, which causes an increase in the blood sugar level.
- (3) Kupffer cells are macrophages that are attached to the luminal surface or inserted in the endothelial lining of hepatic sinusoids. In this site, Kupffer cells play a key role in host defense by removing foreign, toxic and infective substances from the portal blood and by releasing beneficial mediators.
- (4) The sebaceous glands are holocrine glands found in the skin of mammals. They secrete an oily substance called sebum (Latin, meaning for or tallow) that is made of fat (lipids) and the debris of dead fat-producing cells. These glands exist in humans throughout the skin except in the palms of the hands and soles of the feet. Sebum acts to protect and waterproof hair and skin, and keep them from becoming dry, brittle, and cracked. It can also inhibit the growth of microorganisms on skin.

165. (2) There are two types of photoreceptors in the human retina, rods and cones.

Rods are responsible for vision at low light levels (scotopic vision). They do not mediate color vision, and have a low spatial acuity.

Comes are active at higher light levels (photopic vision), are capable of color vision and are responsible for high spatial acuity. The central fovea is populated exclusively by cones. There are 3 types of cones which we will refer to as the short-wavelength sensitive cones, the middle-wavelength sensitive cones and the long-wavelength sensitive cones or S-cone, M-cones, and L-cones for short.

The light levels where both are operational are called mesopic

- 166. (1). Polygenic inheritance is a pattern responsible for many features that seem simple on the surface. Many traits such as height, shape, weight, color, and metabolic rate are governed by the cumulative effects of many genes. Polygenic traits are not expressed as absolute or discrete characters, as was the case with Mendel's pea plant traits. Instead, polygenic traits are recognizable by their expression as a gradation of small differences (a continuous variation).
- 167. (2) A test cross determines whether a particular characteristic of a plant or animal is homozygous dominant (pure bred) or heterozygous dominant (hybrid)

When you look at someone with a dominant trait, you can't tell whether they are homozygous or heterozygous. For instance, if someone has brown eyes, they could be BB or Bb. Mendel always started his crosses with a purebred (homozygous) P generation. How did he know his dominant parent was homozygous? He developed what is known as a test cross. He took a recessive (in our case blue eyes, or bb) and mated it with a pea plant showing the dominant trait. If in the F₁ generation the offspring were all dominant, then the dominant parent was probably BB. If there were any recessives in the F₁, then the dominant parent had to be Bb.

168. (3) Renain is a coagulating enzyme occurring in the gastric juice of the calf, forming the active principle of rennet and able to curdle milk.

Helicase is an enzyme that is capable of unwinding the double helix structure of DNA.

Hyaluronidase: An enzyme that catalyzes the breakdown of hyaluronic acid in the body, thereby increasing tissue permeability to fluids. Also called spreading factor.

By catalyzing the hydrolysis of hyaluronic acid, a major constituent of the interstitial barrier, hyaluronidase lowers the viscosity of hyaluronic acid, thereby increasing tissue permeability. It is, therefore, used in medicine in conjunction with other drugs in order to speed their dispersion and delivery. The most common application is in ophthalmic surgery, in which it is used in combination with local anesthetics. It also increases the absorption rate of parenteral fluids given by hypodermodysis, and is an adjunct in subcutaneous urography for improving resorption of radiopaque agents.

 Lyelne is an essential amino acid, which means that it is essential to human health but cannot be manufactured by the body. For this reason, lysine must be obtained from food.

Glycine is a non-essential, neutral, genetically coded amino acid. It is the only protein-forming amino acid without a center of chirality

Thiamine is one of the B vitamins, a group of water-soluble vitamins that perticipate in many of the chemical reactions in the body.

(2) Myocin is one of the major proteins of muscle. Myosin can slide along filaments of actin, generating force as it does so and using ATP as its motive energy source.

Oxytocin is a hormone produced by the pituitary gland that stimulates contractions of the uterus during labor and release of milk during breast-feeding.

Gastrin is a polypeptide hormone secreted by the mucous lining of the stornach; induces the secretion of gastric juice

(4) Sensory nerve is a nerve that passes impulser from receptors toward or to the central nervous system. Optic nerve is the largest sensory nerve of the eye; carries impulses for sight from the retina to the brain

Oculomotor is a motor nerve and is the third of twelve paired cranial nerves. It controls most of the eye movements (cranical nerves IV and VI also do some), constriction of the pupil, and holding the eyelid open

Vague is a mixed nerve that supplies the pharynx and larynx and lungs and heart and esophagus and stomach and most of the abdominal viscera.

169. (4) Human leg bones: trochanter, fernur, petella, tibia, fibula, tarsals, metatarsals & phalanges.

170. (3) Haemophilia: A disease in which the blood fails to clot. The most common form, primarily effecting males, is caused by a mutation in a gene coding for a clotting protein (factor VIII) inherited as an X-linked recessive phenotype. Red green colour blindness is also a sex linked recessive disease (recessive gene on X chromosome).

171. (4) Chameleone (family Chamaeleonidae) an squamates that belong to one of the best known families. They are known for their ability to change their color, their elongated sticky tongue, and for their eyes which can be moved independently deach other. The name "chameleon" means "eart" lion" and is derived from the Greek words "chame (on the ground, on the earth) and "leon" (lion).

172. (1) Enzymes are catalysts. Most are proteins. Exzymes bind temporarily to one or more of the reactants of the reaction they catalyze. In doing so, the lower the amount of activation energy needed an thus speed up the reaction. In the given grapicurve a shouws normal enzyme reaction, curve!

shows competitive inhibition and curve c shows non-competitive inhibition.

173. (2) Best way to make use of biodegradable pollutants is to use them for producing biogas.

174. (1) Borderline personality disorder - It is a type of psychological disorders in which person is emotionally unstable. They have little sense of self since they feel empty.

A mood disorder is a condition whereby the prevailing emotional mood is distorted or inappropri-

ate to the circumstances.

The two major types of mood disorders are depression (or unipolar depression) and bipolar disorder. Depression (or unipolar depression), including subtypes:

Major Depression

Major Depression (Recurrent)

Major Depression with psychotic symptoms (psychotic depression)

Dysthymia

Postpartum depression

Bipolar disorder, a mood disorder described by alternating periods of mania and depression (and in some cases rapid cycling, mixed states, and psychotic symptoms symptoms). Subtypes include:

Schizophrenia is a mental disorder. It difficult for a person to tell the difference between real and unreal experiences, to think logically, to have normal emotional responses to others, and to behave normally in social situations.

175. (1) Mitochondria have their own genetic material however, and, in sexually reproducing organisms, are inherited only via the cytoplasm of the egg cell

176. (3) These are simple columnar epithelial cells, but in addition, they posses fine hair-like outgrowths, cilia on their free surfaces. These cilia are capable of rapid, rhythmic, wavelike beatings in a certain direction. This movement of the cilia in a certain direction causes the mucus, which is secreted by the goblet cells, to move (flow or stream) in that direction. Ciliated epithelium is usually found in the air passages like the nose. It is also found in the uterus and Fallopian tubes of females. The movement of the cilia propel the ovum to the uterus.

177. (2) Tadpoles treated with thyroxine (thyroid hormone) or iodine will develop at an increased rate, whereas tadpoles treated with thiourea, a thyroid-hormone inhibitor, will metamorphose at a slower rate. Furthermore, a higher concentration of thyroxine will be more effective at speeding metamorphosis than a lower concentration.

Xenopus loevis tadpoles that arrest development and remain as larvae for several years sometimes occur spontaneously in laboratory populations. These tadpoles cease development at an early hindlimb stage, but continue to grow and develop into grossly deformed giants. Giant tadpoles lack thyroid glands, and differ in morphology and behaviour from normal larvae. They are negatively buoyant, typically with small and partially solidified lungs, and have greatly enlarged fat bodies. Giant tadpoles have mature gonads with eggs and sperm, whereas normal tadpoles of the same stage have undifferentiated gonads. Larval reproduction has never been reported in anurans, but gonadal development decoupled from metamorphosis brings these giants the closest of any anurans to being truly neotenic. We discuss behavioural and morphological factors that may hinder both reproduction in giant Xenopus larvae and the evolution of neoteny in anurans in general. Experimental treatment with exogenous thyroid hormone induces some, but not complete, metamorphic changes in these giants. The limbs and head progress through metamorphosis; however, all tadpoles die at the stage when the tail would normally be resorbed. The disproportionate growth of tissues and organs in giant tadpoles may preclude complete metamorphosis, even under exogenous thyroid hormone induction.

178. (1) Biliharzia is a human disease caused by parastic worms called Schistosomes. Over one billion humans are at risk worldwide and approximately 300 millions are infected. Bilharzia is common in the tropics where ponds, streams and irrigation canals harbor bilharzia-transmitting snails. Parasite larvee develop in snails from which they infect humans, their definitive host, in which they mature and reproduce.

Leptospirosis is a potentially serious illness that can affect many parts of the body. Leptospirosis is caused by Leptospira interrogans, a corkscrewshaped bacterium (spirochete). Leptospirosis-causing bacteria are common worldwide, especially in tropical countries with heavy rainfall. Infected rodents and other wild and domestic animals pass the bacteria in their urine. The bacteria can live for a long time in fresh water, damp soil, vegetation, and mud. Flooding after heavy rainfall helps spread the bacteria in the environment. People get leptospirosis by contact with fresh water, damp soil, or vegetation contaminated by the urine of infected animals. People who cance, raft, wade, or swim in contaminated lakes, rivers, and streams can get leptospirosis. Leptospirosis is also a problem for people who work in contaminated flood plains or wet agricultural settings. Leptospirosis bacteria can enter the body through broken skin and mucousmembranes. The bacteria can also enter the body when a person swallows contaminated food or water, including water swallowed during water sports. Once in the bloodstream, the bacteria can

reach all parts of the body and cause signs and symptoms of illness

Hepatitis A is an inflammation of the liver caused by a virus, the hepatitis A virus (HAV). It varies in severity, running an acute course, generally starting within two to six weeks after contact with the virus, and lasting no longer than two or three months. HAV may occur in single cases after contact with an infected relative or sex partner. Alternately, epidemics may develop when food or drinking water is contaminated by the feces of an infected person.

Diarrhoea: Diarrhoea is a symptom of infection caused by a host of bacterial, viral and parasitic organisms most of which can be spread by contaminated water. It is more common when there is a shortage of clean water for drinking, cooking and cleaning and basic hygiene is important in prevention. Water contaminated with human faeces for example from municipal sewage, septic tanks and latrines is of special concern. Animal faeces also contain microorganisms that can cause diarrhoea. Diarrhoea can also spread from person to person, accravated by poor personal hygiene. Food is another major cause of diarrhoea when it is prepared or stored in unhugienic conditions. Water can contaminate food during imigation, and fish and seafood from polluted water may also contribute to the disease.

Guinea worm infection: Infection occurs when the parasitic guinea worm resides within the body. Infection is not apparent until a pregnant female worm prepares to expel embryos. The infection is rarely fatal, but the latter stage is painful. The infection is also referred to as dracunculiasis, and less commonly as drecontiasis.

Fflieriesis: The thread-like, parasitic filarial worms Wuchereria bancrofti and Brugia malayi that cause lymphatic filariasis live almost exclusively in humans. These worms lodge in the lymphatic system, the network of nodes and vessels that maintain the delicate fluid balance between the tissues and blood and are an essential component for the body's immune defence system. They live for 4-6 years, producing millions of immature microfilariae (minute larvae) that circulate in the blood.

The disease is transmitted by mosquitoes that bite infected humans and pick up the microfilariae that develop, inside the mosquito, into the infective stage in a process that usually takes 7-21 days. The larvee, then migrate to the mosquitoes' biting mouth-parts, ready to enter the punctured skin following the mosquito bite, thus completing the cycle.

Amoebic dysentery (amoebiasis) is an infection of the intestine (gut) caused by an amoeba called Entamoeba histolytica, which, among other things, can cause. Amoebae are parasites that are found

in contaminated food or drink. They enter the body through the mouth when the contaminated food or drink is swallowed. The amoebae are then able to move through the digestive system and take up residence in the intestine and cause an infection. Amoebic dysentery is passed on by careless or negligent hygiene where contaminated food and drink is consumed without adequate heat treatment.

Salads washed with contaminated water are a common method of spread.

179. (2) The Maximum Contaminant Level (MCL) of nitrate as nitrogen (NO3-N) at 10 mg/L (or 10 parts per million) for the safety of drinking water. Nitrate levels at or above this level have been known to cause a potentially fatal blood disorder in infants under six months of age called methemoglobinemia or "blue-baby" syndrome; in which there is a reduction in the oxygen-carrying capacity of blood.

Pneumoconlosis is a lung condition that is caused by inhaling particles of mineral dust, usually while working in a high-risk, mineral-related industry. At first, irritating mineral dust can trigger lung inflammation, which causes areas of the lung to be temporarily damaged. Over time, these areas can progress to form tough, fibrous tissue deposits. This stage of pneumoconiosis is called fibrosis. Fibrosis stiffens the lungs and interferes with the lung's normal exchange of oxygen and carbon dioxide. Coal worker's pneumoconiosis is a respiratory disease caused by inhaling coal dust for prolonged periods (Black lung disease)

Non-Hodgidn's lymphoma is cancer that originates in lymphatic system, the disease-fighting network spread throughout the body. non-Hodokin's lymphoma, tumors develop from white blood cells (lymphocytes). These tumors can occur at different locations in the body. There are more than 30 types of non-Hodgkin's lymphoma. Non-Hodgkin's lymphoma is more than seven times as common as the other general type of lymphoma - Hodgkin's disease Causes. Normally, white blood cells (lymphocytes) go through a predictable life cycle. Old lymphocytes die, and body creates new ones to replace them. But in non-Hodgkin's lymphoma, body produces abnormal lymphocytes that continue to divide and grow uncontrollably. This oversupply of lymphocytes crowds into lymph nodes, causing them to swell. Doctors don't know what exactly causes non-Hodgkin's lymphoma. But researchers believe that activation of certain abnormal genes may be involved in the development of all cancers, including lymphomas.

Methane being a green house gas has direct effect on ozone depletion (cause of skin cancers) Exposure to asbestos, a group of minerals found in housing and industrial building materials can cause a variety of medical problems, such as Mesothelioma.

Studies have shown that people who are exposed to high amount of benzene are at risk for cancer. Benzene is a chemical found in gasoline, smoking, and pollution. Often called the "asbestos cancer", Mesothelioma is a disease affecting the abdomen, chest and areas around the heart

180. (2) A coronary anglogram is a special type of X-ray which looks at the blood vessels of the heart. If you have angina or have had a recent heart attack you may be asked to undergo this test. Normai X-rays would not show up blood vessels. A type of due called 'contrast' is injected into the blood vessels of the heart that allows us to see them clearly. The catheter (a thin long length of plastic tubing), which is used to deliver the contrast 'dye' to the heart, is inserted via a large blood vessel in the groin. You will have this area cleaned and a local anaesthetic drug will be injected into the skin so that you do not feel any pain. Once this drug has been injected you will feel pressure and 'tugging' at the site but no pain. The catheter is passed through a large blood vessel in the groin or arm all the way up to the coronary arteries. You may be asked to hold your breath or cough during this procedure. Once the operator enters the blood vessels of the heart a small amount of contrast will be injected so that the blood vessels show up on the X-rays. Rarely this may cause some discomfort in the chest.

181. (3) BRINJAL (Solanaum melongena) is susceptible to nematode invasion due to multiple cropping pattern, good moisture level prevailing and continued presence of host. Rootknot caused by meloidogure incognita is the most destructive nematode.

Potato late blight is one of the most devastating plant diseases. Potato late blight, caused by Phytophthora infestars

Citrus canker is a highly contagious disease of citrus crops caused by the bacterium Xanthomonas axonopodis pathovar citri.

Pigeon pea cyst nematode -Heterodera cajani

182. (4) The basic chromosome number of wheat is 7 (x = 7) and its hexaploid species contains 42 (6 x 7) chromosomes. Thus, it's monosomic (one chromosome rhissing) contains 41 (42 = 1) chromosomes, haploid 21 (42/2) chromosomes. Nullisomic (one chromosome pair missing) contains 40 (40 = 2) chromosomes and trisomic contains (42 + 1) chromosomes.

183. (2). Grafting is a method of plant propagation widely used in horticulture, where the tissues of one plant are encouraged to fuse with those of another. It is most commonly used for the propagation of trees and shrubs grown commercially. (Grafting is limited to dicots and gymnosperms. Monocots lack the vascular cambium required.)

In most cases, one plant is selected for its roots, and this is called the stock or rootstock. The other plant is selected for its stems, leaves, flowers, or fruits and is called the **action**.

In stern grafting, a common grafting method, a shoot of a selected, desired plant cultivar is grafted onto the stock of another type. In another common form called budding, a dormant side bud is grafted on the stern of another stock plant, and when it has fused successfully, it is encouraged to grow by cutting out the stern above the new bud.

For successful grafting to take place, the vascular cambium tissues of the stock and scion plants must be placed in contact with each other. Both tissues must be kept alive till the graft has taken, usually a period of a few weeks. Successful grafting only requires that a vascular connection takes place between the two tissues. A physical weak point often still occurs at the graft, because the structural tissue of the two distinct plants, such as wood may not fuse.

184. (4) Evert concluded that the most likely function of P-protein is to seal the sieve plate pores of injured sieve elements as a rapid first line of defense.

185. (2) Myxomycetes also called Mycetozoa, phylum of funguslike organisms within the kingdom Protista, commonly known as true stime molds. They exhibit characteristics of both protozoans (one-celled microorganisms) and fungi. Distributed worldwide, they usually occur in decaying plant material.

186. (4) Benthan and Hooker did not know the affinities of the families placed under series viii. Ordines anomaly and the families were tentatively grouped together.

1. POLYPETALAE

Series i. THALAMIFLORAE

Series ii. DISCIFLORAE

Series iii. CALYCIFLORAE

GAMOPETALAE Series i. INFERAE

Series ii. HETEROMERAE

Series iii. BICARPELLATAE

3. MONOCHLAMYDEA E

Series i. CURVEMBRYEAE

Series ii. MULTIOVULATAE AQUATICAE

Series III. MULTIOVULATAE TERRESTRES

Series iv. MICREMBRYEAE

Series v. DAPHNALES

Series vi. ACHLAMYDOSPOREAE

Series vii. UNISEXUALES

Series viii. ORDINES ANOMALI

- 187. (3) Hiradin: An anticoagulant ("bloodthinner"). Hirudin is the active principle in the salivary secretion of leeches. The name hirudin is from Hirudo medicinalis, the name of the medicinal leech. In 1884 John Haycraft in Strasbourg found that leeches contained a substance with anticoagulant properties. This anticoagulant in leech saliva was isolated in the 1950s and found to be an antithrombin (an inhibitor of thrombin). The primary chemical structure of hirudin was determined in 1976. Hirudin is produced from transgenic Brassica napus.
- 188. (2) The graft consists of a top portion (the scion), which is the desired plant, growing on the roots of another plant (the stock). The quality of flowers and fruits of such a composite plant produced is mainly determined by scion. (also refer Ans. 183)
- 189. (4) Present in photosynthetic bacterian and blue green algae, chromatophore is a simple body that lacks the complex internal structure and chemically simpler than chlorophyll of plants.
- 190. (4) In fungi, a large sporophore, or fruiting body, in which sexually produced spores are formed on the surface of club-shaped structures (basidia). Basidiocarps are found among the members of the class Basidiomycetes (q.u.), with the exception of the rust and smut fungi. The largest basidiocarps include giant puffballs (Calvatia gigantea), which can be 1.6 m (5.25 feet) long, 1.35 m broad
- 191. (4) Leghaemoglobin: Form of haemoglobin found in the nitrogen-fixing root-nodules of legumes. Binds oxygen, and thus protects the nitrogen-fixing enzyme, nitrogenase, that is oxygen sensitive.
- 192. (1) The Avena geo-curvature test is a bioassay for auxin-type growth regulators. Measurement of the effect of a known or suspected biologically active substance on living material is called as bioassay. Avena curvature test was first time performed by Went for measuring the activity fo auxin hormone.
- 193. (4) New Plastids arise from preexisting plastids by a fission like process.
- 194. (3) Salvia, Calotropis and Mussanda are insect pollinating flowers Kadam and Kigelia are bat pollinating flowers

Triticum, Zea mays, Cannabis and Pinus are wind pollinating flowers.

195. (3) Montreal Protocol (16 september 1987) 27 industrialised countries agreed to limit production of chlorofluorocarbons to half the level of 1986.

- Montreal protocol limits the production and use of ozone depleting substances such as CFCs. To date more than 175 countries have signed the montreal protocol.
- 196. (4) A keystone species is a species whose very presence contributes to a diversity of life and whose extinction would consequently lead to the extinction of other forms of life. Keystone species help to support the ecosystem (entire community of life) of which they are a part.
- 197. (3) Genetic diversity is a characteristic of ecosystems and gene pools that describes an attribute which is commonly held to be advantageous for survival that there are many different versions of otherwise similar organisms. For example, the Irish potato famine can be attributed in part to the fact that there were so few different genetic strains of potatoes in the country, making it easier for one virus to infect and kill much of the crop.
- 198. (3) RNA synthesis continues until RNA polymerase reaches a site on the DNA called the terminator or non-sence codons. Non-sense codons signal the end of protein molecule's synthesis (UAA, UAG, UGA)
- 199. (2) A biosphere reserve is a large area protected for its 'natural' beauty this may include the various plants, the animals feeding on the plants and the primary hunters who in turn feed on the herbivores (although man is a part of the biosphere none have been housed in reserves till date!). National parks are much the same except for the fact that they are usually smaller in size. The origin of the word sanctuary comes from the word sanctuarium (Latin origin). It means "a place of safety for injured or hurt animals". Here the animals recoup their numbers away from their "inhuman human enemies
- 200. (1) Somaclonal variation is the term used to describe the variation seen in plants that have been produced by plant tissue culture. Chromosomal rearrangements are an important source of this variation. Somaclonal variation is not restricted to, but is particularly common in plants regenerated from callus. The variations can be genotypic or phenotypic, which in the later case can be either genetic or epigenetic in origin. Typical genetic alterations are: changes in chromosome numbers (polyploidy) and aneuploidy), chromosome structure (translocations, deletions and duplications) and DNA sequence (base mutations). Typical epigenetic related events are: gene amplification and gene methylation.