

DR ACADEMY

DO RIGHT FOR GENUINE EDUCATION

KCET EXAMINATION PHYSICS – 2019


HELD ON 30-04-2019 (TIME : 10.30 AM TO 11.50 AM)

1. A particle which are initially at rest move towards each other under the action of their mutual attraction. If their speeds are v and $2v$ at any instant, then the speed of centre of mass of the system is

1) $2v$ 2) zero 3) $1.5v$ 4) v

ANS : 2

2. A particle is moving uniformly along a straight line as shown in the figure. During the motion of the particle from A to B, the angular momentum of the particle about 'O'



- 1) Increases
2) Decreases
3) Remains constant
4) First increases then decreases

ANS : 3

3. A satellite is orbiting close to the earth and has a kinetic energy K . The minimum extra kinetic energy required by it to just overcome the gravitation pull of the earth is

1) K 2) $2K$ 3) $\sqrt{3}K$ 4) $2\sqrt{2}K$


ANS : 1

4. A wire is stretched such that its volume remains constant. The poisson's ratio of the material of the wire is

1) 0.50 2) -0.50 3) 0.25 4) -0.25

ANS : 1


5. A cylindrical container containing water has a small hole at height of $H=8\text{cm}$ from the bottom and at a depth of 2cm from the top surface of the liquid. The maximum horizontal distance travelled by the water before it hits the ground (x) is



1) 8 cm 2) $4\sqrt{2}$ cm 3) 4 cm 4) 6 cm

ANS : 1

6. A transparent medium shown relation between i and r as shown. If the speed of light in vacuum is c the Brewster angle for the medium is



1) 30° 2) 45° 3) 60° 4) 90°

ANS : 3

7. In Young's double slit experiment, using monochromatic light of wavelength λ , the intensity of light at a point on the screen where path difference is λ is K units. The intensity of light at a point where path difference is $\frac{\lambda}{3}$ is

1) K 2) $\frac{K}{4}$ 3) $4K$ 4) $2K$

ANS : 2

8. Due to Doppler's effect the shift in wavelength observed is 0.1\AA for a star producing wavelength 6000\AA . Velocity of recession of the star will be

- 1) 25 km/s 2) 10 km/s
3) 5 km/s 4) 20 km/s

ANS : 3

9. An electron is moving with an initial velocity $\vec{V} = V_0 \hat{i}$ and is in a uniform magnetic field $\vec{B} = B_0 \hat{j}$. Then its de Broglie wavelength

- 1) Remains constant
2) Increases with time
3) Decreases with time
4) Increases and decreases periodically

ANS : 1

10. Light of certain frequency and intensity incident on a photosensitive material causes photoelectric effect. If both the frequency and intensity are doubled, the photoelectric saturation current becomes

- 1) Quadrupled 2) Doubled
3) Halved 4) Unchanged

ANS : 2

11. In a cyclotron a charged particle
- 1) Undergoes acceleration all the time
 - 2) Speeds up between the dees because of the magnetic field
 - 3) Speeds up in de
 - 4) Slows down within a dee and speeds up between dees

ANS : 1

12. The numbers of turns in a coil of Galvanometer is tripled, then

- 1) Voltage sensitivity increase 3 times and current sensitivity remains constant
- 2) voltage senility remains constant and current sensitivity increases 3 times
- 3) Both voltage and current sensitivity remains constant
- 4) Both voltage sand current sensitivity decreases by 33%

ANS : 4

13. A circular loop of magnetic moment M is in an arbitrary orientation in an external uniform magnetic field \vec{B} . The work done to rotate the loop by 30° about an axis perpendicular to its plane is

- 1) MB 2) $\sqrt{3} \frac{MB}{2}$ 3) $\frac{MB}{2}$ 4) Zero

ANS : 4

14. In a permanent magnet at room temperature

- 1) Magnetic moment of each molecule is zero
- 2) The individual molecules has non zero magnetic moment which are all perfectly aligned
- 3) Domains are partially aligned
- 4) Domains are all perfectly aligned


ANS : 3

15. Coercivity of a magnet where the ferromagnetic gets completely demagnetized is $3 \times 10^3 \text{ Am}^{-1}$. The minimum current reburied to be passed in a solenoid having 1000 turns per meter, so that the magnet gets completely demagnetized when placed inside the solenoid is

- 1) 20 mA 2) 60 mA 3) 3A 4) 6A

ANS : 3

16. Which of the following nuclei has shorter mean life?



- 1) A 2) B 3) C 4) all

ANS : 1

17. The conductivity of semiconductor increases with increase in temperature because

- 1) Number density of charge carriers increases
- 2) relaxation time increases
- 3) Both number density of charge carriers and relaxation time increases
- 4) Number density of current carriers increases, relaxation time decreases but effect of decrease in relaxation time is much less than increase in number density


ANS : 4

18. For a transistor amplifier, the voltage gain

- 1) Remains constant for all frequencies
- 2) Is high at high and low frequencies and constant in the middle frequency range
- 3) Is low at high and low frequencies and constant and mid frequencies
- 4) Constant at high frequencies and low at low frequencies

ANS : 3

19. In the following circuit, what are P and Q?



- 1) P=0, Q=0
- 2) P=1, Q=0
- 3) P=0, Q=1
- 4) P=1, Q=1

ANS : 3

20. An antenna uses electromagnetic waves of frequency 5MHz. For proper working the size of the antenna should be

- 1) 15m
- 2) 300m
- 3) 15km
- 4) 3km

ANS : 1

21. A magnetic needle has a magnetic moment of $5 \times 10^{-2} \text{ Am}^2$ and moment of inertia $8 \times 10^{-6} \text{ kgm}^2$. It has a period of oscillation of 2s in a magnetic field \vec{B} . The magnitude of magnetic field is approximately

- 1) $1.6 \times 10^{-4} \text{ T}$
- 2) $0.4 \times 10^{-4} \text{ T}$
- 3) $3.2 \times 10^{-4} \text{ T}$
- 4) $0.8 \times 10^{-4} \text{ T}$


ANS : GRACE ($15.7 \times 10^{-4} \text{ T}$)

22. A toroid has 500 turns per metre length. If it carries a current of 2A, the magnetic energy density inside the toroid is

- 1) 0.628 J/m^3
- 2) 0.314 J/m^3
- 3) 6.28 J/m^3
- 4) 3.14 J/m^3

ANS : 1

23. Consider the situation given in figure. The wire AB is slide on the fixed rails with a constant velocity. If the wire AB is replaced by a semi-circular wire, the magnitude of the induced current will



- 1) Increase
- 2) Remain same
- 3) Decrease
- 4) Increase or decrease depending on whether the semicircle bulges towards the resistance or away from it


ANS : 2

24. The frequency of an alternating current is 50 Hz. What is the minimum time taken by current to reach its peak value from rms value?

- 1) $5 \times 10^{-3} \text{ s}$
- 2) $2.5 \times 10^{-3} \text{ s}$
- 3) 0.02 s
- 4) $10 \times 10^{-3} \text{ s}$

ANS : 2

25. The readings of ammeter and voltmeter in the following circuit are respectively



- 1) 1.2A, 120 V
- 2) 1.5 A, 100 V
- 3) 2.7 A, 220 V
- 4) 2.2 A, 220 V

ANS : 4

26. A certain charge 2Q is divided at first into two parts q_1 and q_2 . Later the charges are placed at a certain distance. If the force of interaction between two charges is maximum then

$$\frac{Q}{q_1} = \underline{\hspace{2cm}}$$

- 1) 4
- 2) 2
- 3) 1
- 4) 0.5

ANS : 3

27. A particle of mass m and charge q is placed at rest in uniform electric field E and then released. The kinetic energy attained by the particle after moving a distance is

1) qEy^2 2) qE^2y 3) qEy 4) q^2Ey


ANS : 3

28. An electric dipole is kept in non-uniform electric field. It generally experiences

1) A force and torque
2) A force but not a torque
3) A torque but not a force
4) Neither a force nor a torque

ANS : 1

29. The figure gives the potential V as a function of distance through four regions on x -axis. Which of the following is true for the magnitude of the electric field E in these regions?



1) $E_A > E_B > E_C > E_D$
2) $E_A = E_C$ and $E_B < E_D$
3) $E_B = E_D$ and $E_A < E_C$
4) $E_A < E_B < E_C < E_D$

ANS : 2

30. A system of two charges separated by a certain distance apart stores electrical potential energy. If the distance between them is increased, the potential energy of the system,

1) Increases in any case
2) Decreases in any case
3) May increase or decrease
4) Remains the same


ANS : 3

31. If P , Q and R are physical quantities having different dimensions, which of the following combinations can never be a meaningful quantity?


1) $\frac{P-Q}{R}$ 2) $PQ-R$ 3) $\frac{PQ}{R}$ 4) $\frac{PR-Q^2}{R}$

ANS : 3

32. The given graph shows the variation of velocity (v) with position (x) for a particle moving along a straight line



Which of the following graph shows the variation of acceleration (a) with position (x)?



ANS : 2

33. The trajectory of a projectile projected from origin is given by the equation $y = x - \frac{2x^2}{5}$. The initial velocity of the projectile is

1) $\frac{2}{5} \text{ ms}^{-1}$ 2) 5 ms^{-1} 3) 25 ms^{-1} 4) $\frac{5}{2} \text{ ms}^{-1}$

ANS : 2

34. An object with mass 5 kg is acted upon by a force $\vec{F} = (-3\hat{i} + 4\hat{j}) \text{ N}$. If its initial velocity at $t = 0$ is $\vec{v} = (6\hat{i} - 12\hat{j}) \text{ ms}^{-1}$, the time at which it will just have a velocity along y -axis is

1) 5 s 2) 10 s 3) 2 s 4) 15 s

ANS : 2

35. During inelastic collision between two objects, which of the following quantity always remains conserved?

- 1) Total kinetic energy
- 2) Total mechanical energy
- 3) Total linear momentum
- 4) Speed of each body

ANS : 3

36. In Rutherford experiment, for head-on collision of α - particles with a gold nucleus, the impact parameter is

- 1) Zero
- 2) of the order of 10^{-14} m
- 3) of the order of 10^{-10} m
- 4) of the order of 10^{-6} m

ANS : 1

37. Frequency of revolution of an electron revolving in n^{th} orbit H-atom is proportional to

- 1) $\frac{1}{n^2}$
- 2) n
- 3) n independent of n
- 4) $\frac{1}{n^3}$

ANS : 4

38. A hydrogen atom in ground state absorbs 10.2 eV of energy. The orbital angular momentum of the electron is increased by

- 1) 1.05×10^{-34} Js
- 2) 2.11×10^{-34} Js
- 3) 3.16×10^{-34} Js
- 4) 4.22×10^{-34} Js

ANS : 1

39. The end product of decay of ${}_{90}\text{Th}^{232}$ is ${}_{82}\text{Pb}^{208}$. The number of α and β particles emitted are respectively

- 1) 3, 3
- 2) 6, 4
- 3) 6, 0
- 4) 4, 6

ANS : 2

40. Two protons are kept at a separation of 10nm. Let F_n and F_e be the nuclear force and the electromagnetic force between them

- 1) $F_e = F_n$
- 2) $F_e \gg F_n$
- 3) $F_e \ll F_n$
- 4) F_e and F_n differ only slightly


ANS : 2

41. Two metal plates are separated by 2cm. The potentials of the plates are -10V and +30V. The electric field between the two plates is

- 1) 500 V/m
- 2) 1000 V/m
- 3) 2000 V/m
- 4) 3000 V/m

ANS : 3

42. The equivalent capacitance between A and B is



- 1) 50 pF
- 2) $\frac{100}{3}$ pF
- 3) 150 pF
- 4) 300 pF

ANS : 2

43. A capacitor of capacitance C charged by an amount Q is connected in parallel with an uncharged capacitor of capacitance $2C$. The final charges on the capacitors are

- 1) $\frac{Q}{2}, \frac{Q}{2}$
- 2) $\frac{Q}{4}, \frac{3Q}{4}$
- 3) $\frac{Q}{3}, \frac{2Q}{3}$
- 4) $\frac{Q}{5}, \frac{4Q}{5}$

ANS : 3

44. Though the electron drift velocity is small and electron charge is very small, a conductor can carry an appreciably large current because

- 1) electron number density is very large
- 2) drift velocity of electron is very large
- 3) electron number density depends on temperature
- 4) relaxation time is small

ANS : 1

45. Masses of three wires of copper are in the ratio 1:3:5 and their lengths are in the ratio 5:3:1. The ratio of their electrical resistance are

- 1) 1:3:5
- 2) 5:3:1
- 3) 1:15:125
- 4) 125:15:1


ANS : 3

46. An aluminium sphere is dipped into water. Which of the following is true ?

- 1) Buoyancy will be less in water at 0°C than that in water at 4°C
- 2) Buoyancy will be more in water at 0°C than that in water at 4°C
- 3) Buoyancy in water at 0°C will be same as that in water at 4°C
- 4) Buoyancy may be more or less in water at 4°C depending on the radius of the sphere

ANS : 1

47. A thermodynamic system undergoes a cyclic process ABC as shown in the diagram. The work done by the system per cycle is



- 1) 750J 2) -1250J 3) -750J 4) 1250J

ANS : 3

48. One mole of O_2 gas is heated at constant pressure starting at $27^\circ C$. How much energy must be added to the gas as heat to double its volume ?

- 1) Zero 2) 450R 3) 750R 4) 1050R

ANS : 4

49. A piston is performing S.H.M. in the vertical direction with a frequency of 0.5Hz. A block of 10kg is placed on the piston. The maximum amplitude of the system such that the block remains in contact with the piston is

- 1) 1m 2) 0.5m 3) 1.5m 4) 0.1m

ANS : 1

50. The equation of a stationary wave is $y = 2 \sin\left(\frac{\pi x}{15}\right) \cos(48\pi t)$. The distance between a node and its next antinode is

- 1) 7.5 units 2) 1.5 units
3) 22.5 units 4) 30 units

ANS : 1

51. An inductor of inductance L and resistor R joined together in series and connected by a source of frequency ω . The power dissipated in the circuit is

- 1) $\frac{R^2 + \omega^2 L^2}{V}$ 2) $\frac{V^2 R}{R^2 + \omega^2 L^2}$
3) $\frac{V}{R^2 + \omega^2 L^2}$ 4) $\frac{V^2 R}{\sqrt{R^2 + \omega^2 L^2}}$

ANS : 2

52. An electromagnetic wave is travelling in x-direction with electric field vector given by $\vec{E}_y = E_0 \sin(kx - \omega t) \hat{j}$. The correct expression for magnetic field vector is

- 1) $\vec{B}_y = E_0 C \sin(kx - \omega t) \hat{j}$
2) $\vec{B}_z = E_0 C \sin(kx - \omega t) \hat{k}$
3) $\vec{B}_y = \frac{E_0}{C} \sin(kx - \omega t) \hat{j}$
4) $\vec{B}_z = \frac{E_0}{C} \sin(kx - \omega t) \hat{k}$


ANS : 4

53. The phenomenon involved in the reflection of radio-waves by ionosphere is similar to

- 1) reflection of light by plane mirror
2) total internal reflection of light in air during a mirage
3) dispersion of light by water molecules during the formation of a rainbow
4) scattering of light by air particles

ANS : 2

54. A point object is moving uniformly towards the pole of a concave mirror of focal length 25cm along its axis as shown below. The speed of the object is 1 ms^{-1} . At $t=0$, the distance of the object from the mirror is 50cm. the average velocity of the image formed by the mirror between time $t=0$ and $t=0.25\text{s}$ is



- 1) 40 cm s^{-1} 2) 20 cm s^{-1}
3) Zero 4) Infinity


ANS : 4

55. A certain prism is found to produce a minimum deviation of 38° . It produces a deviation of 44° when the angle of incidence is either 42° or 62° . What is the angle of incidence when it is undergoing minimum deviation ?

- 1) 30° 2) 40° 3) 49° 4) 60°

ANS : 3

56. In the given circuit, the current through 2Ω resistor is



- 1) 0.2A 2) 0.3A 3) 0.4A 4) 0.1A


ANS : 3

57. Kirchhoffs junction rule is a reflection of

- 1) Conservation of current density vector
2) Conservation of energy
3) Conservation of momentum
4) Conservation of charges

ANS : 4

58. The variation of terminal potential difference (V) with current flowing through a cell is as shown



The emf and internal resistance of the cell are

- 1) 3V, 2Ω 2) 3V, 0.5Ω
3) 6V, 2Ω 4) 6V, 0.5Ω


ANS : 2

59. In a potentiometer experiment, the balancing point with a cell is at a length 240cm. On shutting the cell with a resistance of 2Ω , the balancing length becomes 120cm. The internal resistance of the cell is

- 1) 4Ω 2) 2Ω 3) 1Ω 4) 0.5Ω

ANS : 2

60. The magnetic field at the centre 'O' in the given figure is



- 1) $\frac{7}{14} \frac{\mu_0 I}{R}$ 2) $\frac{5}{12} \frac{\mu_0 I}{R}$ 3) $\frac{3}{10} \frac{\mu_0 I}{R}$ 4) $\frac{\mu_0 I}{12R}$

ANS : 2