

**M.Phil. DEGREE EXAMINATION, DECEMBER 2010****Mathematics****COMMUTATIVE ALGEBRA**

(CBCS—2008 onwards)

Time : 3 Hours

Maximum : 75 Marks

Answer **all** questions.  $(5 \times 15 = 75)$ 

1. (a) Define the term exact sequence of R-modules.  
Prove that :

(i) for any exact sequence  
 $O \rightarrow N' \xrightarrow{f} N \xrightarrow{g} N'' \rightarrow O$  of R-modules, the  
 induced sequence :

$O \rightarrow \text{Hom}_R(M_1 N') \xrightarrow{f^*} \text{Hom}_R(M_1 N) \xrightarrow{g^*} \text{Hom}_R(M_1 N'') \rightarrow O$  is exact  
 where M is a -Rmodule.

- (ii) An R-module P is projective if and only if for any surjective homomorphism  $g : M \rightarrow M''$  the induced homomorphism  $g^* : \text{Hom}_R(P, M) \rightarrow \text{Hom}_R(P, M'')$  is surjective.

(Or)

(b) For an  $R$ -module  $M$  show that the following are equivalent :

(i) A sequence  $0 \rightarrow N' \xrightarrow{f} N \xrightarrow{g} N'' \rightarrow 0$  of  $R$ -modules is exact if and only if the tensored sequence :

$$0 \rightarrow M \otimes N' \xrightarrow{f} M \otimes N \xrightarrow{g} M \otimes N'' \rightarrow 0 \text{ is exact.}$$

(ii)  $M$  is  $R$ -flat and for any  $R$ -module  $N$ ,

$$M \otimes_l N = 0 \text{ implies } N = 0.$$

(iii)  $M$  is  $R$ -flat and for any  $R$ -homomorphism

$$f : N' \rightarrow N, \quad \text{the induced map}$$

$$f^* : M \otimes N' \rightarrow M \otimes N \text{ is zero implies that}$$

$$f = 0.$$

Prove that  $M$  is faithfully flat if and only if  $M$  is flat and for each maximal ideal  $m$  of  $R$ ,  $m^M \neq M$ .

2. (a) Prove :

- (i) Let  $R$  be a local ring. Any finitely generated projective  $R$ -module is free.
- (ii) Let  $R$  be a local ring with maximal ideal  $m$  and  $M$  be a finitely presented  $R$ -module. If the canonical map  $u_M : m \otimes_R M \rightarrow M$  given by  $u_M : (a \otimes x) = ax, a \in m, x \in M$  is injective then  $M$  is free.

(Or)

- (b) For a ring  $R$ , describe  $R_S$ , for a multiplicatively closed subset  $S$  of  $R$ . If  $S$  is a multiplicatively closed set and  $f : R \rightarrow R_S$  be the natural map given by  $f(a) = (a/1)$ , show that (i) every ideal of  $R_S$  is an extended ideal (ii) the prime ideals of  $R_S$  are in 1 - 1 correspondence with the prime ideals of  $R$  not intersecting  $S$  ; iii)  $f$  preserves the ideal operations of taking finite sums, products, intersections and radical.

3. (a) Let  $R$  be an Artinian ring. Show that (i) every prime ideal of  $R$  is maximal (ii) there are finitely many maximal ideals of  $R$  (iii) the nil radical is nilpotent.

(Or)

- (b) Define the term “length of a module”. State and prove the Jordan-Hölder theorem w.r.t. composition series and length.
4. (a) Prove :
- (i) Let  $R$  and  $S$  be domains and  $S$  integral over  $R$ . The  $R$  is a field if and only if  $S$  is a field.
- (ii) Let  $R, S$  be as in (i) for any prime ideal  $P$  of  $R$ , there exists a prime ideal  $P'$  of  $S$  such that  $P' \cap R = P$ .

- (iii) Let  $S$  be an integral extension of  $R$ ,  $f: R \rightarrow \Omega$  be a ring homomorphism of  $R$  into an algebraically closed field  $\Omega$ . Then  $f$  can be extended to a ring homomorphism  $g: S \rightarrow \Omega$ .

(Or)

- (b) Let  $R$  be an integrally closed domain with quotient field  $K$  and  $S$  normal extension of  $R$  with Galois group  $G = G(S/K)$ . Show that :

- (i)  $G$  is the group of  $R$ -automorphisms of  $S$ .
- (ii) Two prime ideals  $P'$  and  $Q'$  of  $S$  lie over the same prime ideal of  $R$  if and only if there exists some  $\sigma \in G$  with  $\sigma(P') = Q'$ .

5. (a) Define the term valuation ring. Let  $p$  be a fixed prime and  $\mathbb{R}\mathbb{C}\mathbb{Q}$ , where  $\mathbb{Q}$  is the field of rationals is defined by :

$$R = \left\{ p^r \frac{m}{n} : r \geq 0, (n, p) = 1, (n, p) > 1 \right\}. \text{ Show that } R$$

is a valuation ring. Prove (i) the ideals of a valuation ring are totally ordered by inclusion (ii) if the ideals of a domain  $V$  in the quotient field  $K$  are totally ordered by inclusion then  $V$  is a valuation ring of  $K$ .

(Or)

- (b) Let  $R$  be a Noetherian local domain, with unique maximal ideal  $m \neq 0$ , and  $K$  be the quotient field of  $R$ . Show that the following are equivalent :

- (i)  $R$  is a discrete valuation ring ;
- (ii)  $R$  is a principal ideal domain ;

- (iii)  $m$  is principal.
- (iv)  $R$  is integrally closed and every non-zero prime ideal of  $R$  is maximal.
- (v) every non-zero ideal of  $R$  is a power of  $m$ .

\_\_\_\_\_ \*\*\* \_\_\_\_\_

**M.Phil. DEGREE EXAMINATION, DECEMBER 2010****Mathematics****FUNCTIONAL ANALYSIS**

(CBCS–2008 onwards)

Time : 3 Hours

Maximum : 75 Marks

Answer **all** questions. (5 × 15 = 75)

1. (a) Let  $\wedge$  be a linear functional on a Topological vector space  $X$ . Assume  $\wedge x \neq 0$  for some  $x \in X$ . Prove that the following four properties are equivalent :

- (i)  $\wedge$  is continuous.
- (ii) The null space  $\mathcal{N}(\wedge)$  is closed.
- (iii)  $\mathcal{N}(\wedge)$  is not dense in  $X$ .
- (iv)  $\wedge$  is bounded in some neighbourhood  $\vee$  of 0.

(Or)

(b) (i) Suppose  $K$  and  $C$  are subsets of a topological vector space  $X$ ,  $K$  is compact,  $C$  is closed and  $K \cap C = \emptyset$ . Prove that  $0$  has a neighbourhood  $V$  such that  $(K + V) \cap (C + V) = \emptyset$ .

(ii) Prove that every locally compact topological vector space  $X$  has finite dimension.

(8 + 7)

2. (a) (i) State and prove Baire's theorem.

(ii) State and prove Banach-Steinhaus theorem.

(7 + 8)

(Or)

(b) State and prove Open mapping theorem.

3. (a) State and prove the Banach-Alaoglu theorem. State and prove any two applications of the above theorem.

(9 + 6)

(Or)

- (b) Let  $M$  be a subspace of a real vector space  $X$ .

Let  $p: X \rightarrow \mathbb{R}$  satisfy  $p(x + y) \leq p(x) + p(y)$  and  $p(tx) = p(x)$  if  $x \in X, y \in X, t \geq 0$ . Suppose  $f: M \rightarrow \mathbb{R}$  is linear and  $f(x) \leq p(x)$  on  $M$ . Prove that there exists a linear  $\wedge: X \rightarrow \mathbb{R}$  such that  $\wedge(x) = f(x), (x \in M)$  and  $-p(-x) \leq \wedge(x) \leq p(x), x \in X$ .

4. (a) (i) Suppose  $X$  and  $Y$  are normed spaces. Prove that to each  $T \in \mathcal{B}(X, Y)$  there corresponds a unique  $T^* \in \mathcal{B}(Y^*, X^*)$  that satisfies  $\langle Tx, y^* \rangle = \langle x, T^* y^* \rangle$  for all  $x \in X$  and all  $y^* \in Y^*$ . Also prove that  $\|T^*\| = \|T\|$ .

- (ii) Suppose  $X$  and  $Y$  are Banach spaces and  $T \in \mathcal{B}(X, Y)$ . Prove that  $T$  is compact if, and only if,  $T^*$  is compact.

(Or)

- (b) Suppose  $X$  is a Banach Space,  $T \in \mathcal{B}(X)$  and  $T$  is compact. Then prove the following :

- (i) If  $\lambda \neq 0$ , then the four numbers  
 $\alpha = \dim \mathcal{N}(T - \lambda I)$ ,  $\beta = \dim (X / \mathcal{R}(T - \lambda I))$ ,  
 $\alpha^* = \dim \mathcal{N}(T^* - \lambda I)$  and  
 $\beta^* = \dim (X^* / \mathcal{R}(T^* - \lambda I))$  are equal and finite.

- (ii) if  $\lambda \neq 0$ , and  $\lambda \in \sigma(T)$  then  $\lambda$  is an eigen value of  $T$  and of  $T^*$ .

- (iii)  $\sigma(T)$  is compact, atmost countable and has atmost one limit point namely 0.

5. (a) State and prove Bishop's theorem. Give an example to illustrate Bishop's theorem.

(Or)

- (b) Let  $G$  be a compact group. Prove that there exists a unique regular Borel probability measure  $m$  which is left invariant, right invariant and satisfies the relation

$$\int_G f(x) dm(x) = \int_G f(x^{-1}) dm(x)$$

where  $f \in C(G)$ .

\_\_\_\_\_ \*\*\* \_\_\_\_\_

**M.Phil. DEGREE EXAMINATION, DECEMBER 2010****Mathematics****MEASURE THEORY**

(CBCS–2008 onwards)

Time : 3 Hours

Maximum : 75 Marks

Answer **all** questions. $(5 \times 15 = 75)$ 

1. (a) (i) Define measurable space and measurable sets.

(5)

- (ii) Suppose  $\mathcal{m}$  is a  $\sigma$ -algebra in  $X$ , and  $Y$  is a topological space let  $f$  map  $X$  into  $Y$

1. If  $\Omega$  is the collection of all sets  $E \subset Y$  such that  $f^{-1}(E) \in \mathcal{m}$ , then  $\Omega$  is a  $\sigma$ -algebra in  $Y$
2. If  $f$  is measurable and  $E$  is a Borel set in  $Y$  then  $f^{-1}(E) \in \mathcal{m}$ .

(10)

(Or)

- (b) Suppose  $f$  and  $g \in L'(\mu)$  and  $\alpha$  and  $\beta$  are complex numbers

then prove that  $\alpha f + \beta g \in L'(\mu)$  and

$$\int_X (\alpha f + \beta g) d\mu = \alpha \int_X f d\mu + \beta \int_X g d\mu .$$

(15)

2. (a) (i) Explain  $C_c(X)$ . (5)

- (ii) State and prove Urysohn's lemma. (10)

(Or)

- (b) (i) Define  $\sigma$ -finite measure. (3)

- (ii) Let  $X$  be a locally compact Hausdorff space in which every open set is  $\sigma$ -compact. Let  $\lambda$  be any positive Borel measure on  $X$  such that  $\lambda(K) < \infty, \forall K$  then prove that  $\lambda$  is regular.

(12)

3. (a) (i) If  $\phi$  is convex on  $(a, b)$  then  $\phi$  is continuous on  $(a, b)$  where  $\phi$  is a real function on  $(a, b)$ .

(5)

- (ii) State and prove Jensen's inequality.

(10)

(Or)

- (b) (i) Prove that  $L^p(\mu)$  is a complete metric space, for  $1 \leq p \leq \infty$  and for every positive measure  $\mu$

(10)

- (ii) If  $\{f_n\}$  is a Cauchy sequence in  $L^p(\mu)$  with limit  $f$  then prove  $\{f_n\}$  has a subsequence which converges pointwise almost everywhere to  $f(x)$ .

(5)

4. (a) (i) Let  $\mu$  be a complex measure on a  $\sigma$ -algebra  $\mathcal{M}$  in  $X$ , then there is a measurable function  $h$  such that  $|h(x)| = 1 \quad \forall x \in X$  such that  $d\mu = h d|\mu|$

(8)

- (ii) State and prove the Hahn-decomposition theorem.

(7)

(Or)

- (b) State and prove the Riesz representation theorem for a unique regular complex Borel measure.

5. (a) (i) Prove that weak  $L^1$  contains  $L^1$

(5)

(ii) If  $f \in L^1(\mathbb{R}^K)$ , then prove that almost every  $x \in \mathbb{R}^K$  is a Lebesgue point of  $f$ .

(10)

(Or)

(b) If  $T(B(x, r))$  is Lebesgue measurable, the set  $V$  is open in  $\mathbb{R}^K$ ,  $T: V \rightarrow \mathbb{R}^K$  is continuous and  $T$  is differentiable at some point  $x \in V$  then prove that.

$$\lim_{r \rightarrow 0} \frac{m(T(B(x, r)))}{m(B(x, r))} = \Delta(T'(x))$$

(15)

\_\_\_\_\_ \*\*\* \_\_\_\_\_